



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Higgs boson production at the LHC using the q_T subtraction formalism at N^3LO QCD

Cieri, Leandro ; Chen, Xuan ; Gehrmann, Thomas ; Glover, E W N ; Huss, Alexander

Abstract: We consider higher-order QCD corrections to Higgs boson production through gluon-gluon fusion in the large top quark mass limit in hadron collisions. We extend the transverse-momentum (q_T) subtraction method to next-to-next-to-next-to-leading order (N³LO) and combine it with the NNLO Higgs-plus-jet calculation to numerically compute differential infrared-safe observables at N³LO for Higgs boson production in gluon fusion. To cancel the infrared divergences, we exploit the universal behaviour of the associated q_T distributions in the small- q_T region. We document all the necessary ingredients of the transverse-momentum subtraction method up to N³LO. The missing third-order collinear functions, which contribute only at $q_T = 0$, are approximated using a prescription which uses the known result for the total Higgs boson cross section at this order. As a first application of the third-order q_T subtraction method, we present the N³LO rapidity distribution of the Higgs boson at the LHC.

DOI: [https://doi.org/10.1007/jhep02\(2019\)096](https://doi.org/10.1007/jhep02(2019)096)

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-170875>

Journal Article

Published Version

Originally published at:

Cieri, Leandro; Chen, Xuan; Gehrmann, Thomas; Glover, E W N; Huss, Alexander (2019). Higgs boson production at the LHC using the q_T subtraction formalism at N^3LO QCD. Journal of High Energy Physics, 2019(2):96.

DOI: [https://doi.org/10.1007/jhep02\(2019\)096](https://doi.org/10.1007/jhep02(2019)096)

RECEIVED: August 17, 2018
 REVISED: December 14, 2018
 ACCEPTED: February 2, 2019
 PUBLISHED: February 14, 2019

Higgs boson production at the LHC using the q_T subtraction formalism at N³LO QCD

Leandro Cieri,^{a,b} Xuan Chen,^b Thomas Gehrmann,^b E.W.N. Glover^c and Alexander Huss^d

^a*INFN, Sezione di Milano-Bicocca,
 Piazza della Scienza 3, I-20126 Milano, Italy*

^b*Physik-Institut, Universität Zürich,
 CH-8057 Zurich, Switzerland*

^c*Institute for Particle Physics Phenomenology, Durham University,
 Durham, DH1 3LE, U.K.*

^d*Theoretical Physics Department, CERN,
 1211 Geneva 23, Switzerland*

E-mail: lcieri@physik.uzh.ch, xuan.chen@uzh.ch,
thomas.gehrmann@uzh.ch, e.w.n.glover@durham.ac.uk,
alexander.huss@cern.ch

ABSTRACT: We consider higher-order QCD corrections to Higgs boson production through gluon-gluon fusion in the large top quark mass limit in hadron collisions. We extend the transverse-momentum (q_T) subtraction method to next-to-next-to-next-to-leading order (N³LO) and combine it with the NNLO Higgs-plus-jet calculation to numerically compute differential infrared-safe observables at N³LO for Higgs boson production in gluon fusion. To cancel the infrared divergences, we exploit the universal behaviour of the associated q_T distributions in the small- q_T region. We document all the necessary ingredients of the transverse-momentum subtraction method up to N³LO. The missing third-order collinear functions, which contribute only at $q_T = 0$, are approximated using a prescription which uses the known result for the total Higgs boson cross section at this order. As a first application of the third-order q_T subtraction method, we present the N³LO rapidity distribution of the Higgs boson at the LHC.

KEYWORDS: QCD Phenomenology

ARXIV EPRINT: [1807.11501](https://arxiv.org/abs/1807.11501)

Contents

1	Introduction	1
2	The q_T subtraction formalism at N³LO	2
3	The Higgs boson total cross section at N³LO	8
3.1	Implementation and setup of the numerical calculations	10
3.2	The numerical extraction of C_{N3}	12
4	The rapidity distribution of the Higgs boson	15
4.1	The NNLO rapidity distribution	15
4.2	Numerical stability of the N ³ LO rapidity distribution	17
4.3	The rapidity distribution of the Higgs boson at N ³ LO	19
5	Conclusions and outlook	20
A	Fixed-order expressions	22
B	Convolutions at N³LO	24

1 Introduction

The most straightforward and successful (as well as systematically improvable) approach to calculations for processes at high-momentum scales M in QCD is a perturbative expansion in the strong coupling $\alpha_s(M^2)$. Cross sections are written as a series expansion in the parameter α_s and an improvement in accuracy is obtained by calculating an increasing number of coefficients in the series. Until a few years ago, the standard for such calculations was next-to-leading order (NLO) accuracy. Recent years have seen a number of next-to-next-to-leading order (NNLO) results for many important processes of interest, such that the emerging standard for precision calculations relevant for LHC phenomenology is the second non-trivial order in the strong coupling α_s .

Reducing the theoretical uncertainties remains one of the main motivations for the extension from NLO to NNLO accuracy. This is particularly relevant in two distinct situations. Firstly, NNLO corrections are mandatory for those processes where NLO corrections are comparable in size to the leading order (LO) contribution, both to establish the convergence of the perturbative expansion and to obtain reliable predictions. Secondly, many benchmark processes demand theoretical predictions with the highest possible precision to be able to fully exploit the extraordinary experimental precision that is achievable for this class of processes. Such “standard candles” are not only indispensable tools in detector calibration but also allow for a precise extraction of Standard Model (SM) parameters and parton distribution functions (PDFs).

Extending the perturbative accuracy of QCD calculations to one order higher implies developing new methods and techniques to achieve the cancellation of infrared (IR) divergences that appear at intermediate steps of the calculations. The past few years have witnessed a great development in NNLO subtraction prescriptions. The transverse momentum (q_T) subtraction method [1–3], the N -jettiness subtraction [4, 5], projection-to-Born [8], residue subtraction [6, 7], and the antenna subtraction method [9–11] have all been successfully applied for LHC phenomenology.

However, in view of the impressive and continuously improving quality of the measurements performed at the LHC, even NNLO accuracy is in some cases not sufficient to match the demands of the LHC data. Typically, these are processes in which the size of the NLO corrections are comparable with the LO, and where the NNLO corrections still exhibit large effects such that the size of the theoretical uncertainties remains larger than the experimental uncertainties.

This motivated a new theoretical effort to go beyond NNLO to include the next perturbative order: the next-to-next-to-next-to-leading order (N^3 LO). Sum rules, branching fractions [12] and deep inelastic structure functions [13] have been known to this order for quite some time. At present, the only hadron collider observables for which N^3 LO QCD corrections have been calculated are the total cross section for Higgs boson production in gluon fusion [14, 15] and in vector boson fusion [16]. First steps have been taken towards more differential observables by computing several N^3 LO threshold expansion terms to the Higgs boson rapidity distribution in gluon fusion [17, 18]. Moreover, the projection-to-Born method has been most recently extended to compute fully differential distributions to N^3 LO, with a proof-of-principle calculation [19] of jet production in deep inelastic scattering.

In this paper we extend the q_T subtraction method at N^3 LO to compute Higgs boson production differentially in the Higgs boson rapidity at N^3 LO accuracy. The paper is organized as follows: in section 2 we recall briefly the main ideas of the q_T subtraction formalism and we present the necessary ingredients up to N^3 LO, specifying which elements are known analytically and identifying the missing coefficients at N^3 LO. In section 3 we present a prescription for approximating the missing collinear functions at N^3 LO based on the unitarity property of the integral of the transverse momentum distribution. In section 4, we apply the q_T subtraction formalism at N^3 LO to produce differential distributions in the rapidity of the Higgs boson. To validate our approach, section 4.1 quantifies the quality of the approximations by repeating them at NNLO, where all of the ingredients to q_T subtraction are known. We assess the magnitude of different sources of systematic uncertainties at N^3 LO in section 4.2, yielding final results for the N^3 LO Higgs boson rapidity distribution and the associated theoretical uncertainty in section 4.3. Finally, in section 5 we summarize our results.

2 The q_T subtraction formalism at N^3 LO

This section is devoted to present briefly the transverse-momentum subtraction formalism to N^3 LO in perturbative QCD. The method is illustrated in its general form and spe-

cial attention is paid to the case of Higgs boson production through gluon-gluon fusion. The q_T subtraction formalism presented in this section is the third-order extension of the subtraction method originally proposed in refs. [1–3].

We consider the inclusive hard scattering reaction

$$h_1(p_1) + h_2(p_2) \rightarrow F(\{q_i\}) + X, \quad (2.1)$$

where h_1 and h_2 denote the two hadrons which collide with momenta p_1 and p_2 producing the identified colourless final-state system F , accompanied by an arbitrary and undetected final state X . The colliding hadrons have centre-of-mass energy \sqrt{s} , and are treated as massless particles

$$s = (p_1 + p_2)^2 = 2p_1 \cdot p_2.$$

The observed final state F consists of a generic system of non-QCD partons composed of *one or more* colour singlet particles (such as vector bosons, photons, Higgs bosons, Drell-Yan (DY) lepton pairs and so forth) with momenta q_i^μ ($i = 3, 4, 5, \dots$). The total momentum of the system F is denoted by

$$q^\mu = \sum_i q_i^\mu,$$

and the kinematics of the system can be expressed in terms of the total invariant mass M ,

$$M^2 = q^2,$$

the transverse momentum q_T with respect to the direction of the colliding hadrons (omitting the azimuthal dependence), and the rapidity in the centre-of-mass system of the hadronic collision, Y ,

$$Y = \frac{1}{2} \ln \left(\frac{p_2 \cdot q}{p_1 \cdot q} \right).$$

The fully differential hadronic cross section can therefore be written as

$$\frac{d\sigma^F}{dq_T^2 dM^2 dY} = \int_0^1 d\xi_1 \int_0^1 d\xi_2 \frac{d\hat{\sigma}_{ab}^F(\xi_1 p_1, \xi_2 p_2)}{dq_T^2 dM^2 dY} f_{a/h_1}(\xi_1, \mu_F) f_{b/h_2}(\xi_2, \mu_F), \quad (2.2)$$

where $d\hat{\sigma}_{ab}$ is the differential partonic cross section, ξ_1, ξ_2 are the partonic momentum fractions and $f_{c/h}$ the distribution function for finding parton c in hadron h . Since F is colourless, the LO partonic cross section can be either initiated by $q\bar{q}$ annihilation, as in the case of the Drell-Yan process, or by gluon-gluon fusion, as in the case of Higgs boson production. In the case of the Born cross section, the kinematics of the colour-neutral system F is fully constrained such that

$$\begin{aligned} \frac{d\hat{\sigma}_{\text{LO};ab}^F(\xi_1 p_1, \xi_2 p_2)}{dq_T^2 dM^2 dY} &= \delta_{ca} \delta_{\bar{c}b} \delta(q_T^2) \delta(M^2 - \xi_1 \xi_2 s) \delta(Y - \ln(\xi_1/\xi_2)/2) d\hat{\sigma}_{\text{LO};c\bar{c}}^F(\xi_1 p_1, \xi_2 p_2) \\ &= \delta_{ca} \delta_{\bar{c}b} \delta(q_T^2) \delta\left(\xi_1 - \frac{M}{\sqrt{s}} e^+ Y\right) \delta\left(\xi_2 - \frac{M}{\sqrt{s}} e^- Y\right) \frac{1}{s} d\hat{\sigma}_{\text{LO};c\bar{c}}^F(\xi_1 p_1, \xi_2 p_2). \end{aligned} \quad (2.3)$$

In order to explain the basic idea of the subtraction formalism, we first notice that at LO, the transverse momentum q_T of the final state system F is identically zero. Therefore, as long as $q_T > 0$, the N^n LO QCD contributions (with $n \geq 1$) are given by the N^{n-1} LO QCD contributions to the $F + \text{jet}(s)$ final state. Consequently, if $q_T > 0$ we have:

$$\left. \frac{d\sigma_{N^n\text{LO}}^F}{dq_T^2 dM^2 dY} \right|_{q_T > 0} \equiv \frac{d\sigma_{N^{n-1}\text{LO}}^{F+\text{jet}(s)}}{dq_T^2 dM^2 dY}, \quad (2.4)$$

where the notation N^n LO stands for: N^0 LO = LO, N^1 LO = NLO, N^2 LO = NNLO and so forth. Equation (2.4) implies that if $q_T > 0$ the infrared (IR) divergences that appear in the computation of $d\sigma_{N^n\text{LO}}^F|_{q_T > 0}$ are those already present in $d\sigma_{N^{n-1}\text{LO}}^{F+\text{jet}(s)}$. Therefore, provided that the IR singularities involved in $d\sigma_{N^{n-1}\text{LO}}^{F+\text{jet}(s)}$ can be handled and cancelled with the available subtraction methods at N^{n-1} LO, the only remaining singularities at N^n LO are associated with the limit $q_T \rightarrow 0$ and we treat them with the q_T subtraction method. Since the small- q_T behaviour of the transverse momentum distribution is well known through the resummation program [20–32] of logarithmically-enhanced contributions to transverse-momentum distributions, we can (in principle) exploit this knowledge to construct the necessary N^n LO counterterms (CT) to subtract the remaining singularity at $q_T = 0$, thereby promoting the q_T subtraction method proposed in ref. [1] to N^n LO.

The generic form of the q_T subtraction method [1] for the N^n LO cross section is

$$\frac{d\sigma_{N^n\text{LO}}^F}{dq_T^2 dM^2 dY} = \mathcal{H}_{N^n\text{LO}}^F \otimes \frac{d\sigma_{\text{LO}}^F}{dq_T^2 dM^2 dY} + \left[\frac{d\sigma_{N^{n-1}\text{LO}}^{F+\text{jet}(s)}}{dq_T^2 dM^2 dY} - \frac{d\sigma_{N^n\text{LO}}^{F\text{CT}}(q_T)}{dq_T^2 dM^2 dY} \right], \quad (2.5)$$

where the symbol “ \otimes ” denotes convolutions over the momentum fractions and the flavour indices of the incoming partons and is explicitly defined as

$$\begin{aligned} \mathcal{G}(\dots) \otimes \frac{d\sigma^F}{d\mathcal{O}} &\equiv \int_0^1 d\xi_1 \int_0^1 d\xi_2 \int_0^1 dz_1 \int_0^1 dz_2 \\ &\times \frac{d\hat{\sigma}_{ab}^F(\xi_1 z_1 p_1, \xi_2 z_2 p_2)}{d\mathcal{O}} \mathcal{G}_{ab \leftarrow cd}(\dots; z_1, z_2) f_{c/h_1}(\xi_1, \mu_F) f_{d/h_2}(\xi_2, \mu_F). \\ &= \int_0^1 dx_1 \int_0^1 dx_2 \int_{x_1}^1 \frac{dz_1}{z_1} \int_{x_2}^1 \frac{dz_2}{z_2} \\ &\times \frac{d\hat{\sigma}_{ab}^F(x_1 p_1, x_2 p_2)}{d\mathcal{O}} \mathcal{G}_{ab \leftarrow cd}(\dots; z_1, z_2) f_{c/h_1}\left(\frac{x_1}{z_1}, \mu_F\right) f_{d/h_2}\left(\frac{x_2}{z_2}, \mu_F\right). \end{aligned} \quad (2.6)$$

The counterterm $d\sigma_{N^n\text{LO}}^{F\text{CT}}$ constitutes the contribution to the N^n LO cross section which cancels the divergences of $d\sigma_{N^{n-1}\text{LO}}^{F+\text{jet}(s)}$ in the limit $q_T \rightarrow 0$ and renders the term in square brackets finite for all values of q_T . The n -th order counterterm can be written as

$$\frac{d\sigma_{N^n\text{LO}}^{F\text{CT}}(q_T)}{dq_T^2 dM^2 dY} = \Sigma_{N^n\text{LO}}^F(q_T) \otimes \frac{d\sigma_{\text{LO}}^F}{dM^2 dY}, \quad (2.7)$$

where we note that the dependence of the function $\Sigma_{N^n\text{LO}}^F(q_T)$ on the transverse momentum q_T is *not* kinematically related to the Born-level process.

The functions $\Sigma_{\text{N}^n\text{LO}}^F(q_T)$ and $\mathcal{H}_{\text{N}^n\text{LO}}^F$ correspond to the n -th order truncation of the perturbative series in α_s of the functions

$$\Sigma_{c\bar{c}\leftarrow ab}^F(q_T; z_1, z_2) = \sum_{n=1}^{\infty} \left(\frac{\alpha_s}{\pi}\right)^n \Sigma_{c\bar{c}\leftarrow ab}^{F;(n)}(q_T; z_1, z_2), \quad (2.8)$$

$$\mathcal{H}_{c\bar{c}\leftarrow ab}^F(z_1, z_2) = \delta_{ca}\delta_{\bar{c}b} \delta(1-z_1)\delta(1-z_2) + \sum_{n=1}^{\infty} \left(\frac{\alpha_s}{\pi}\right)^n \mathcal{H}_{c\bar{c}\leftarrow ab}^{F;(n)}(z_1, z_2), \quad (2.9)$$

where the labels a and b stand for the partonic channels of the N^nLO correction that are mapped to that the Born cross section. The function $\Sigma^F(q_T)$ embodies all the terms of the form $\log(q_T^2/M^2)$ that are divergent in the limit $q_T \rightarrow 0$ and reproduces the logarithmically singular behaviour of $d\sigma^{F+\text{jet(s)}}$ in the small- q_T limit. Terms proportional to $\delta(q_T^2)$ as well as IR finite terms are absorbed in the perturbative factor \mathcal{H}^F . The hard coefficient function $\mathcal{H}_{\text{N}^n\text{LO}}^F$ thus encodes all the IR finite terms of the n -loop contributions.

According to the transverse momentum resummation formula [2] and using the Fourier transformation between the conjugate variables q_T and the impact parameter b , the perturbative hard function \mathcal{H}^F and the corresponding counterterm are obtained by the fixed-order truncation of the identity

$$\begin{aligned} \left(\Sigma^F(q_T) + \mathcal{H}^F \delta(q_T^2)\right) \otimes \frac{d\sigma_{\text{LO}}^F}{dM^2 dY} &= \frac{1}{s} \int_0^\infty db \frac{b}{2} J_0(bq_T) d\hat{\sigma}_{\text{LO};c\bar{c}}^F(x_1 p_1, x_2 p_2) S_c(M, b) \\ &\times \int_{x_1}^1 \frac{dz_1}{z_1} \int_{x_2}^1 \frac{dz_2}{z_2} [H^F C_1 C_2]_{c\bar{c};ab} f_{a/h_1}\left(\frac{x_1}{z_1}, \frac{b_0^2}{b^2}\right) f_{b/h_2}\left(\frac{x_2}{z_2}, \frac{b_0^2}{b^2}\right), \end{aligned} \quad (2.10)$$

where $b_0 = 2e^{-\gamma_E}$ ($\gamma_E = 0.5772\dots$ is the Euler-Mascheroni constant).

The large logarithmic corrections are exponentiated in the Sudakov form factor $S_c(M, b)$ of the quark ($c = q, \bar{q}$) or of the gluon ($c = g$), which has the following expression:

$$S_c(M, b) = \exp \left\{ - \int_{b_0^2/b^2}^{M^2} \frac{dq^2}{q^2} \left[A_c(\alpha_s(q^2)) \ln \frac{M^2}{q^2} + B_c(\alpha_s(q^2)) \right] \right\}, \quad (2.11)$$

where the functions A and B permit a perturbative expansion in α_s :

$$A_c(\alpha_s) = \sum_{n=1}^{\infty} \left(\frac{\alpha_s}{\pi}\right)^n A_c^{(n)}, \quad B_c(\alpha_s) = \sum_{n=1}^{\infty} \left(\frac{\alpha_s}{\pi}\right)^n B_c^{(n)}. \quad (2.12)$$

Explicit expressions for the coefficients $A_g^{(n)}$ and $B_g^{(n)}$ that are relevant for Higgs production are collected in appendix A up to $n = 3$. In particular, we also give the $B_g^{(3)}$ coefficient in the hard resummation scheme as needed to evaluate eq. (2.10) for $F = H$ at N^3LO .

The analytical form of the function $\Sigma^{F;(3)}$ in eq. (2.8) can be obtained by expanding eq. (2.10) to the corresponding matching order. The full analytical formula for Σ^F is resummation scheme independent order by order in the strong coupling constant. Therefore, the logarithmic singular behaviour for Σ^F at $q_T \rightarrow 0$ at each given order in α_s does not depend on the resummation scheme, and can be validated against the behaviour of the

fixed-order results at small q_T . To fully account for the logarithmically enhanced terms at a given order requires a sufficient depth in the resummation accuracy prior to its fixed-order expansion in eq. (2.8). Specifically, the LO Higgs boson q_T distribution receives singular contributions from up to NLL (next-to-leading-logarithm) resummation [30, 48], the NLO Higgs boson q_T distribution requires the expansion of NNLL resummation [31, 38, 49–51], and the NNLO Higgs boson q_T distribution has been recently validated against the singular contributions from N³LL resummation [52, 53].

The structure of the symbolic factor denoted by $[H^F C_1 C_2]_{c\bar{c};ab}$ in eq. (2.10), depends on the initial-state channel of the Born subprocess and is explained in detail in refs. [34, 35]. Here we limit ourselves to the case in which the final state system F is composed of a single Higgs boson, $F \equiv H$, in which case,

$$[H^H C_1 C_2]_{gg;ab} = H_g^H(\alpha_s(M^2)) \left[C_{ga}(z_1; \alpha_s(b_0^2/b^2)) C_{gb}(z_2; \alpha_s(b_0^2/b^2)) + G_{ga}(z_1; \alpha_s(b_0^2/b^2)) G_{gb}(z_2; \alpha_s(b_0^2/b^2)) \right], \quad (2.13)$$

where H_g^H is the hard-virtual function and respectively C_{ga} and G_{ga} the gluonic helicity-preserving and helicity-flipping hard-collinear coefficient functions.

The gluonic hard-collinear coefficient function $C_{ga}(z; \alpha_s)$ ($a = q, \bar{q}, g$) has the following perturbative expansion

$$C_{ga}(z; \alpha_s) = \delta_{ga} \delta(1-z) + \sum_{n=1}^{\infty} \left(\frac{\alpha_s}{\pi} \right)^n C_{ga}^{(n)}(z). \quad (2.14)$$

In contrast, the perturbative expansion of the helicity flip hard-collinear coefficient function G_{ga} , which is specific to gluon-initiated processes, starts only at $\mathcal{O}(\alpha_s)$, and can be expanded as [34, 35]

$$G_{ga}(z; \alpha_s) = \sum_{n=1}^{\infty} \left(\frac{\alpha_s}{\pi} \right)^n G_{ga}^{(n)}(z). \quad (2.15)$$

The IR finite contribution of the n -loop correction terms to the Born subprocess is contained in the hard-virtual function (which does not depend on z_1 or z_2),

$$H_g^H(\alpha_s) = 1 + \sum_{n=1}^{\infty} \left(\frac{\alpha_s}{\pi} \right)^n H_g^{H;(n)}. \quad (2.16)$$

Using eqs. (2.10) and (2.13), then, after integration over b and dropping the renormalisation group predictable terms that are produced by evolving α_s to a common scale (i.e. setting $\mu_F = \mu_R = M$), we obtain the resummation scheme independent

$$\mathcal{H}_{gg \leftarrow ab}^H(z_1, z_2; \mu_F = \mu_R = M) \equiv H_g^H(\alpha_s) \left[C_{ga}(z_1; \alpha_s) C_{gb}(z_2; \alpha_s) + G_{ga}(z_1; \alpha_s) G_{gb}(z_2; \alpha_s) \right]. \quad (2.17)$$

Note that in the literature, it is often the rapidity-integrated variant $\mathcal{H}_{gg \leftarrow ab}^H(z)$ that is quoted which is related to $\mathcal{H}_{gg \leftarrow ab}^H(z_1, z_2)$ via the convolution

$$\mathcal{H}_{gg \leftarrow ab}^H(z) \equiv \int_0^1 dz_1 \int_0^1 dz_2 \delta(z - z_1 z_2) \mathcal{H}_{gg \leftarrow ab}^H(z_1, z_2). \quad (2.18)$$

The \mathcal{H}^H function in eq. (2.17) can be expanded perturbatively without approximation to any order in the strong coupling constant α_s . Inserting the expansions of the hard functions into eq. (2.17), then,

$$\begin{aligned} \mathcal{H}_{gg \leftarrow ab}^{H;(1)}(z_1, z_2; \mu_F = \mu_R = M) &= \delta_{ga} \delta_{gb} \delta(1 - z_1) \delta(1 - z_2) H_g^{H;(1)} \\ &+ \delta_{ga} \delta(1 - z_1) C_{gb}^{(1)}(z_2) + \delta_{gb} \delta(1 - z_2) C_{ga}^{(1)}(z_1), \end{aligned} \quad (2.19)$$

$$\begin{aligned} \mathcal{H}_{gg \leftarrow ab}^{H;(2)}(z_1, z_2; \mu_F = \mu_R = M) &= \delta_{ga} \delta_{gb} \delta(1 - z_1) \delta(1 - z_2) H_g^{H;(2)} \\ &+ \delta_{ga} \delta(1 - z_1) C_{gb}^{(2)}(z_2) + \delta_{gb} \delta(1 - z_2) C_{ga}^{(2)}(z_1) \\ &+ H_g^{H;(1)} \left(\delta_{ga} \delta(1 - z_1) C_{gb}^{(1)}(z_2) + \delta_{gb} \delta(1 - z_2) C_{ga}^{(1)}(z_1) \right) \\ &+ C_{ga}^{(1)}(z_1) C_{gb}^{(1)}(z_2) + G_{ga}^{(1)}(z_1) G_{gb}^{(1)}(z_2). \end{aligned} \quad (2.20)$$

Explicit expressions for the known fixed-order coefficients are collected in appendix A.

The new third-order contribution is given by

$$\begin{aligned} \mathcal{H}_{gg \leftarrow ab}^{H;(3)}(z_1, z_2; \mu_F = \mu_R = M) &= \delta_{ga} \delta_{gb} \delta(1 - z_1) \delta(1 - z_2) H_g^{H;(3)} \\ &+ \delta_{ga} \delta(1 - z_1) C_{gb}^{(3)}(z_2) + \delta_{gb} \delta(1 - z_2) C_{ga}^{(3)}(z_1) \\ &+ G_{ga}^{(1)}(z_1) G_{gb}^{(2)}(z_2) + G_{ga}^{(2)}(z_1) G_{gb}^{(1)}(z_2) \\ &+ H_g^{H;(1)} \left(\delta_{ga} \delta(1 - z_1) C_{gb}^{(2)}(z_2) + \delta_{gb} \delta(1 - z_2) C_{ga}^{(2)}(z_1) \right) \\ &+ H_g^{H;(2)} \left(\delta_{ga} \delta(1 - z_1) C_{gb}^{(1)}(z_2) + \delta_{gb} \delta(1 - z_2) C_{ga}^{(1)}(z_1) \right) \\ &+ H_g^{H;(1)} C_{ga}^{(1)}(z_1) C_{gb}^{(1)}(z_2) + H_g^{H;(1)} G_{ga}^{(1)}(z_1) G_{gb}^{(1)}(z_2) \\ &+ C_{ga}^{(1)}(z_1) C_{gb}^{(2)}(z_2) + C_{ga}^{(2)}(z_1) C_{gb}^{(1)}(z_2). \end{aligned} \quad (2.21)$$

The second-order helicity-flip functions $G_{ga}^{(2)}(z)$, the third-order collinear functions $C_{ga}^{(3)}(z)$ and the third-order hard-virtual coefficient $H_g^{H;(3)}$ are only known in parts or not at all, thereby presenting an obstacle to applying the q_T subtraction formalism at N³LO. Nevertheless, within the q_T subtraction formalism, all these resummation coefficients can be inferred for any hard scattering process whose corresponding total cross section is known at N³LO. This point is discussed in detail in section 3.

Although the hard-virtual coefficient $H_g^{H;(3)}$ is currently not known in analytical form, parts of it can be inferred from known results in threshold resummation. This relies on the knowledge of the general structure of H_c^F (to all orders), which relates $H_c^{F;(n)}$ to the finite part of the n -loop virtual Matrix Element [35]. To this end, we split $H_g^{H;(3)}$ into two pieces,

$$H_g^{H;(3)} \equiv \tilde{H}_g^{H;(3)} + [H_g^{H;(3)}]_{(\delta_{(2)}^{q_T})}, \quad (2.22)$$

where $\tilde{H}_g^{H;(3)}$ can be computed using the corresponding hard-virtual factor $C_{gg \rightarrow H}^{\text{th}(3)}$ [57] from threshold resummation (in the large- m_t limit) and the exponential equation that relates

hard-virtual coefficients in threshold- and q_T -resummation (eq. (81) of ref. [35]). We find,

$$\begin{aligned}
 \tilde{H}_g^{H;(3)} = & C_A^3 \left(-\frac{15649\zeta_3}{432} - \frac{121\pi^2\zeta_3}{432} + \frac{3\zeta_3^2}{2} + \frac{869\zeta_5}{144} + \frac{215131}{5184} + \frac{16151\pi^2}{7776} - \frac{961\pi^4}{15552} \right. \\
 & \left. + \frac{\pi^6}{810} + \frac{105}{32}\zeta_6 \right) + C_A^2 \left(\frac{605\zeta_3}{72} + \frac{55\pi^2\zeta_3}{36} + \frac{737\pi^2}{432} + \frac{167\pi^4}{432} + \frac{\pi^6}{72} \right) \\
 & + C_A \left(\frac{19\pi^2 L_t}{48} - \frac{55\pi^2\zeta_3}{8} - \frac{\pi^6}{480} + \frac{133\pi^4}{72} + \frac{11399\pi^2}{864} + \frac{63}{32}\zeta_6 \right) \\
 & + N_f^2 \left(\frac{43C_A\zeta_3}{108} - \frac{19\pi^4 C_A}{3240} - \frac{133\pi^2 C_A}{1944} + \frac{2515C_A}{1728} - \frac{7C_F\zeta_3}{6} \right. \\
 & \left. + \frac{4481C_F}{2592} - \frac{\pi^4 C_F}{3240} - \frac{23\pi^2 C_F}{432} \right) \\
 & + N_f \left(\frac{101C_A^2\zeta_5}{72} - \frac{97}{216}\pi^2 C_A^2\zeta_3 + \frac{29C_A^2\zeta_3}{8} + \frac{1849\pi^4 C_A^2}{38880} - \frac{35\pi^2 C_A^2}{243} - \frac{98059C_A^2}{5184} \right. \\
 & + \frac{5C_A C_F\zeta_5}{2} + \frac{13C_A C_F\zeta_3}{2} + \frac{1}{2}\pi^2 C_A C_F\zeta_3 - \frac{63991C_A C_F}{5184} + \frac{11\pi^4 C_A C_F}{6480} \\
 & - \frac{71}{216}\pi^2 C_A C_F + \frac{1}{9}\pi^2 C_A L_t - \frac{5}{36}\pi^2 C_A\zeta_3 - \frac{55C_A\zeta_3}{36} - \frac{5\pi^4 C_A}{54} - \frac{1409\pi^2 C_A}{864} \\
 & \left. - 5C_F^2\zeta_5 + \frac{37C_F^2\zeta_3}{12} + \frac{19C_F^2}{18} \right), \tag{2.23}
 \end{aligned}$$

with $L_t = \ln(M^2/m_t^2)$ and ζ_n denoting the Riemann zeta-function for integer values n ($\zeta_2 = \pi^2/6$, $\zeta_3 = 1.202\dots$, $\zeta_4 = \pi^4/90$). Note that we neglect all the third-order terms in the exponent of eq. (81) in ref. [35], considering the entire $\mathcal{O}(\alpha_s^3)$ correction (in the exponent) as unknown. However, the full top-mass dependence of $H_g^{H;(3)}$ is already fully embodied in $\tilde{H}_g^{H;(3)}$. The currently unknown $[H_g^{H;(3)}]_{(\delta_{(2)}^{q_T})}$ represents a *single* coefficient (of *soft* origin) belonging to the finite part of the structure of the IR singularities contained in the third-order virtual amplitude of the corresponding partonic subprocess $gg \rightarrow H$.

As a consequence, the only missing ingredients to $\mathcal{H}^{H;(3)}$ are the functions $G_{ga}^{(2)}(z)$, $C_{ga}^{(3)}(z)$ and $[H_g^{H;(3)}]_{(\delta_{(2)}^{q_T})}$. The details on their numerical extraction will be discussed in the following section.

3 The Higgs boson total cross section at N³LO

We start this section by reviewing some properties of the hard-scattering function $\mathcal{H}_{c\bar{c} \leftarrow ab}^F$. This function is resummation-scheme independent, but it depends on the specific hard-scattering subprocess $c + \bar{c} \rightarrow F$. The coefficients $\mathcal{H}_{c\bar{c} \leftarrow ab}^{F;(n)}$ of the perturbative expansion in eq. (2.9) can be determined by performing a perturbative calculation of the q_T distribution in the limit $q_T \rightarrow 0$. In the right-hand side of eq. (2.10), the function \mathcal{H}^F controls the strict perturbative normalization of the corresponding total cross section (i.e. the integral of the total q_T distribution). This unitarity-related property can be exploited to determine the coefficients $\mathcal{H}_{c\bar{c} \leftarrow ab}^{F;(n)}$ from the perturbative calculation of the inclusive cross section. In particular, the integral of the full q_T spectrum in eq. (2.5) must reproduce the inclusive

cross section $\sigma^{F(\text{tot.})}$,

$$\sigma_{\text{N}^n\text{LO}}^{F(\text{tot.})} = \int_0^\infty dq_T^2 \frac{d\sigma_{\text{N}^n\text{LO}}^F}{dq_T^2}, \quad \frac{d\sigma_{\text{N}^n\text{LO}}^F}{dq_T^2} \equiv \int dM^2 dY \frac{d\sigma_{\text{N}^n\text{LO}}^F}{dq_T^2 dM^2 dY}. \quad (3.1)$$

Since the hard-scattering function $\mathcal{H}_{c\bar{c} \leftarrow ab}^F$ is accompanied by $\delta(q_T^2)$, we evaluate the q_T spectrum on right-hand side of eq. (2.5) according to the following decomposition [2]

$$\sigma_{\text{N}^n\text{LO}}^{F(\text{tot.})} = \mathcal{H}_{\text{N}^n\text{LO}}^F \otimes \sigma_{\text{LO}}^F + \int_0^\infty dq_T^2 \frac{d\sigma_{\text{N}^n\text{LO}}^{F(\text{fin.})}}{dq_T^2}, \quad (3.2)$$

where $d\sigma^{F(\text{fin.})}$ is directly related to the quantity in square brackets in the right-hand side of eq. (2.5)

$$\frac{d\sigma_{\text{N}^n\text{LO}}^{F(\text{fin.})}}{dq_T^2} \equiv \left[\frac{d\sigma_{\text{N}^{n-1}\text{LO}}^{F+\text{jet(s)}}}{dq_T^2} - \frac{d\sigma_{\text{N}^n\text{LO}}^{F\text{CT}}}{dq_T^2} \right]. \quad (3.3)$$

The relation in eq. (3.2) is valid order-by-order in QCD perturbation theory [2]. If the perturbative coefficients of the fixed-order expansion of $\sigma^{F(\text{tot.})}$, \mathcal{H}^F and $d\sigma^{F(\text{fin.})}/dq_T^2$ are all known, the relation (3.2) has to be regarded as an identity, which can be explicitly checked. Since the fixed-order truncation of $d\sigma^{F(\text{fin.})}/dq_T^2$ is free of any contribution proportional to $\delta(q_T^2)$, its NLO contribution does not contain the coefficient $\mathcal{H}^{F;(1)}$, and so forth. Therefore, $\mathcal{H}^{F;(3)}$ can be isolated from the N^3LO term in eq. (3.2):

$$\left[\sigma_{\text{N}^3\text{LO}}^{F(\text{tot.})} - \sigma_{\text{N}^2\text{LO}}^{F(\text{tot.})} \right] - \int_0^\infty dq_T^2 \left[\frac{d\sigma_{\text{N}^3\text{LO}}^{F(\text{fin.})}}{dq_T^2} - \frac{d\sigma_{\text{N}^2\text{LO}}^{F(\text{fin.})}}{dq_T^2} \right] = \left(\frac{\alpha_s}{\pi} \right)^3 \mathcal{H}^{F;(3)} \otimes \sigma_{\text{LO}}^F, \quad (3.4)$$

where $\alpha_s = \alpha_s(\mu_R^2)$.

If all the components on the left-hand side of eq. (3.4) are known analytically (as it was the case at NNLO in refs. [39, 40]), the function \mathcal{H}^F can be extracted exactly in analytical form. At NLO the extraction of the function $\mathcal{H}^{F;(1)}$ is straightforward for Drell-Yan and Higgs boson production. The function $\mathcal{H}^{F;(2)}$ at NNLO (for Higgs ($F = H$) boson production [39] and Drell-Yan ($F = DY$) [40]) can be obtained with a dedicated analytical computation using the analogue of eq. (3.4) at NNLO. Since the transverse momentum distributions for H +jet and DY +jet at NNLO are not known analytically, eq. (3.4) can be used only numerically to compute $\mathcal{H}^{F;(3)}$.

As was elaborated on at the end of the previous section, the general structure of the coefficient $\mathcal{H}^{F;(3)}$ is not known in analytic form for any hard-scattering process. Nonetheless, within the q_T subtraction formalism, $\mathcal{H}^{F;(3)}$ can be reliably approximated for any hard-scattering process whose corresponding total cross section is known at N^3LO . As identified in eqs. (2.21) and (2.22), the only missing ingredients to $\mathcal{H}^{F;(3)}$ are the functions $G_{ga}^{(2)}(z)$, $C_{ga}^{(3)}(z)$ and $[H_g^{H;(3)}]_{(\delta_{(2)}^{q_T})}$. Their contribution to eq. (3.4) can be approximated as follows:

$$\begin{aligned} & \left[H_g^{H;(3)} \right]_{\delta_{(2)}^{q_T}} \delta_{ga} \delta(1-z_1) \delta_{gb} \delta(1-z_2) \\ & + C_{ga}^{(3)}(z_1) \delta_{gb} \delta(1-z_2) + \delta_{ga} \delta(1-z_1) C_{gb}^{(3)}(z_2) + G_{ga}^{(2)}(z_1) G_{gb}^{(1)}(z_2) + G_{ga}^{(1)}(z_1) G_{gb}^{(2)}(z_2) \\ & \approx C_{N3} \delta_{ga} \delta(1-z_1) \delta_{gb} \delta(1-z_2), \end{aligned} \quad (3.5)$$

where the third-order coefficient C_{N3} embodies the numerical extraction of the hard-virtual coefficient $[H_g^{H;(3)}]_{(\delta_{(2)}^{q_T})}$ *plus* the approximation of the z_i -dependent functions by a numerical constant proportional to $\delta(1 - z_i)$. The resulting coefficient $[H_g^{H;(3)}]_{(\delta_{(2)}^{q_T})}$ is exact since C_{N3} is proportional to $\delta(1 - z)$ (or equivalently $\delta(1 - z_1)\delta(1 - z_2)$). In other words, the approximation that is made in eq. (3.5) is related only to the functions $G_{ga}^{(2)}(z)$ and $C_{ga}^{(3)}(z)$, whose functional dependence on the variable z goes beyond terms proportional to $\delta(1 - z)$, and which involves not only gluon-to-gluon transitions ($a = g$), but also contributions from other parton species ($a = q, \bar{q}$). The latter are not explicitly distinguished in the above approximation, which fully attributes their numerical contribution to the gluon-induced processes.

The method outlined in eq. (3.5) to approximate the unknown terms in the hard-virtual function $\mathcal{H}_{gg \leftarrow ab}^H$ numerically is not new. It was first used in ref. [2] in order to compute the second order function $\mathcal{H}_{gg \leftarrow ab}^{H;(2)}$ numerically at NNLO, providing a reasonable estimate of the exact result to better than 1% accuracy. Notice that eq. (3.5) ensures that one recovers the total cross section (at N³LO in this case) with no approximation. After integration over the transverse momentum q_T , eq. (3.1) provides the same total integral (numerically in this case) as in the fully analytical case. Even more, for IR-safe observables (at fixed order) where the *back-to-back* kinematical configuration ($q_T = 0$) is located at a single phase space point (e.g. the q_T distribution, the angular separation $\Delta\varphi_{\gamma\gamma}$ between the two photons for a Higgs boson decaying into diphotons, etc.), the fixed order result is also exact, i.e. the integral of the analytical unknown terms in eq. (3.5) (which all have $q_T = 0$) is located in one single point of the exclusive differential distributions.

The previous considerations about the approximation underpinning eq. (3.5) were regarding the total cross section or differential distributions in which the Born-like configurations belong to one single phase space point. In order to quantify the quality of the approximation proposed in eq. (3.5) at the differential level when the Born differential cross section populates the entire differential range, we perform a detailed numerical study of the Higgs boson rapidity $Y \equiv y_H$ distribution in section 4.1 at NNLO. Anticipating these results, we find that in the rapidity range $0 \leq y_H \leq 4$ the approximated NNLO result differs by less than 0.2% from the exact NNLO Higgs boson rapidity distribution.

3.1 Implementation and setup of the numerical calculations

To extract the value of C_{N3} , we first introduce the numerical tools and the calculational setup in this section. We use the same setup for the inclusive and differential predictions presented in sections 3.2, 4.1, 4.2 and 4.3.

We consider Higgs boson production in proton-proton collisions at a centre-of-mass energy of $\sqrt{s} = 13$ TeV. In our computation, we set the Higgs boson mass to $M \equiv M_H = 125$ GeV and the vacuum expectation value to $v = 246.2$ GeV. The Born process is initiated via gluon-gluon fusion mediated through a top-quark loop, which can be integrated out in the large- m_t limit ($m_t \rightarrow \infty$). In this limit, the production of the Higgs boson is described through an effective gluon-gluon-Higgs boson vertex [59–61]. The mass of the top quark is taken as $m_t = 173.2$ GeV, which enters in the contributions that have a residual

m_t dependence (e.g. eqs. (A.10) and (2.23) and effective vertex coefficient corrections at N³LO). With the top quark loop replaced by an effective vertex, we consider a five-flavour scheme QCD with all light quarks being massless. We use the central set of the PDF4LHC15 PDFs [62] as implemented in the LHAPDF framework [63] and the associated strong coupling constant with $\alpha_s(M_Z) = 0.118$. Note that we systematically employ the same order in the PDFs (in particular the set PDF4LHC15_nnlo_mc) for the LO, NLO, NNLO and N³LO results presented in this paper. The central factorization and renormalization scale is chosen as $\mu \equiv \mu_R = \mu_F = M_H/2$. The theoretical uncertainty is estimated by varying the default scale choice independently for μ_R and μ_F by factors of $\{1/2, 2\}$ while omitting combinations with $\mu_R/\mu_F = 4$ or $1/4$, resulting in the common seven-point variation of scale combinations.

As stated in section 3 and in ref. [1], the computation of the total cross section or differential distributions with the q_T subtraction formalism can be separated into two main parts by inserting eq. (3.3) into eq. (3.2):

$$\sigma_{\text{N}^n\text{LO}}^{F(\text{tot.})} = \left[\mathcal{H}_{\text{N}^n\text{LO}}^F \otimes \sigma_{\text{LO}}^F - \int_0^\infty dq_T^2 \frac{d\sigma_{\text{N}^n\text{LO}}^{F\text{CT}}}{dq_T^2} \right] + \int_0^\infty dq_T^2 \frac{d\sigma_{\text{N}^{n-1}\text{LO}}^{F+\text{jet}(s)}}{dq_T^2}. \quad (3.6)$$

The contribution $d\sigma^{F+\text{jet}(s)}$ in eq. (3.6) is computed with the parton-level event generator NNLOJET which provides the necessary infrastructure for the antenna subtraction method up to NNLO [9–11]. Processes at NNLO with the structure of $d\sigma^{F+\text{jet}(s)}$ implemented in NNLOJET are: $F = H$ [64], $F = \gamma^*$, Z [65, 66] and $F = W^\pm$ [67]. In this paper we focus on Higgs production $F = H$, where the relevant matrix elements in NNLOJET are: $(H+1)$ -parton production at two loops [68], $(H+2)$ -parton production at one loop [69–71] and $(H+3)$ -parton production at tree-level [72–74]. The subtraction formalism that we are applying to Higgs boson production could be easily extended to Z and W^\pm production [75].

The terms in square brackets in eq. (3.6) for $F = H$ are encoded in a new Monte Carlo generator HN3LO [76] up to the third order in the strong coupling constant. After expanding eq. (2.10) to this order, several non-trivial convolutions emerge and we briefly document the corresponding formulae implemented in HN3LO in appendix B. All our results up to the NNLO level are in full agreement with the Monte Carlo generator HNNLO [1] at the per mille level of accuracy. On the left-hand side of eq. (3.4), the Higgs boson total cross sections at NNLO ($\sigma_{\text{NNLO}}^{H(\text{tot.})}$) and N³LO ($\sigma_{\text{N}^3\text{LO}}^{H(\text{tot.})}$) are also required. We use the analytical coefficient function for the total Higgs boson cross section that was recently calculated in ref. [15] and which is available within the public program ihixs 2 [77]. This program is further used to compute any of the analytical total cross-section ingredients required to extract the coefficient C_{N^3} .

The numerical computation of the integral of the difference $d\sigma_{\text{NNLO}}^{F+\text{jet}(s)} - d\sigma_{\text{N}^3\text{LO}}^{F\text{CT}}$ in eq. (3.3), although finite, requires the introduction of a suitable technical lower bound or q_T^{cut} , since both terms in this difference are logarithmically divergent at $q_T^{\text{cut}} \rightarrow 0$. This technical cut introduces systematic uncertainties to both $d\sigma_{\text{NNLO}}^{F+\text{jet}(s)}$ and $d\sigma_{\text{N}^3\text{LO}}^{\text{CT}}$. Once cancellations between the terms on the right-hand side of eq. (3.6) take place, the numerically calculated total cross sections and differential distributions have to be q_T^{cut} independent




Figure 1. The q_T integrated finite contribution to the cross section of eq. (3.3) at N³LO-only (i.e. N³LO – NNLO) between q_T^{cut} and ∞ , for three different scales ($\mu = \mu_R = \mu_F$).

(within the statistical errors) over some range of q_T^{cut} . At the lower end of this range, numerical instabilities in $d\sigma_{\text{NNLO}}^{F+\text{jet}(s)}$ (arising from the large dynamical range in this calculation) will limit the accuracy of the result, while at the higher end of the range, missing non-logarithmic terms in $d\sigma_{\text{N}^3\text{LO}}^{F\text{CT}}$ will start to become significant. The numerical stability of $d\sigma_{\text{NNLO}}^{F+\text{jet}(s)}$ at small q_T using NNLOJET has been systematically validated for Higgs boson production (with $q_T^{\text{cut}} = 0.7$ GeV in ref. [52]) and Drell-Yan production (with $q_T^{\text{cut}} = 2$ GeV in ref. [53]) at the LHC. In sections 3.2, 4.1, 4.2 and 4.3, we document numerical results obtained with the q_T subtraction formalism using $q_T^{\text{cut}} = (2 \pm 1)$ GeV.

3.2 The numerical extraction of C_{N3}

In the following, we describe the numerical results regarding the extraction of the C_{N3} coefficient and the corresponding N³LO total cross section.

In figure 1 we display the $\sigma_{\text{N}^3\text{LO}}^{H(\text{fin.})}$ at N³LO-only *coefficient* as a function of the q_T^{cut} , i.e., the difference $\sigma_{\text{N}^3\text{LO}}^{H(\text{fin.})} - \sigma_{\text{NNLO}}^{H(\text{fin.})}$. The error bars denote the numerical integration errors from NNLOJET. Since the figure displays cumulant cross sections as function of the lower integration boundary, the central values and errors are fully correlated among the points. Using eq. (2.21) with eq. (3.4) and the value of the resulting integral $\sigma^{H(\text{fin.})}(q_T^{\text{cut}} = 1 \text{ GeV})$ in figure 1, it is possible to obtain the q_T -integrated cross section of the unknown terms on the left-hand side of eq. (3.5) and consequently extract C_{N3} .

The behaviour of $\sigma_{\text{N}^3\text{LO}}^{H(\text{fin.})}$ as a function of q_T^{cut} is shown in figure 1 and gives an estimate of the systematical uncertainty corresponding to the use of this technical cut which turns out to be at the *per mille* level in the domain $q_T^{\text{cut}} = (2 \pm 1)$ GeV for the total Higgs boson cross section at N³LO. More specifically, variations of the q_T^{cut} parameter from $q_T^{\text{cut}} = 1$ GeV to 3 GeV produce variations in the central value of the N³LO contribution to $\sigma^{H(\text{fin.})}$ cross




Figure 2. The numerically extracted C_{N3} coefficient (for three different values of q_T^{cut}) as a function of the combination of scales, as enumerated in table 1. The error bars for each particular C_{N3} point are obtained propagating the statistical uncertainties of the different terms involved in the computation. The red band corresponds to our best estimation for C_{N3} obtained with the central scale $\mu = M_H/2$ at $q_T^{\text{cut}} = 1$ GeV, as detailed in the text.

n	$[\tilde{\mu}_R, \tilde{\mu}_F] \times M_H$	$C_{N3} (q_T^{\text{cut}} = 1 \text{ GeV})$	$C_{N3} (q_T^{\text{cut}} = 2 \text{ GeV})$	$C_{N3} (q_T^{\text{cut}} = 3 \text{ GeV})$
(1)	$[1/2, 1/2]$	-943 ± 222	-967 ± 179	-988 ± 164
(2)	$[1, 1]$	-971 ± 207	-965 ± 168	-989 ± 151
(3)	$[1/4, 1/4]$	-883 ± 243	-866 ± 198	-850 ± 162
(4)	$[1/2, 1]$	-986 ± 222	-1021 ± 179	-1033 ± 179
(5)	$[1, 1/2]$	-990 ± 206	-976 ± 167	-968 ± 158
(6)	$[1/2, 1/4]$	-985 ± 221	-978 ± 181	-923 ± 152
(7)	$[1/4, 1/2]$	-977 ± 243	-859 ± 199	-883 ± 179

Table 1. Extracted values of the C_{N3} coefficients as a function of the q_T^{cut} as shown in figure 2 for each scale choice. In bold typeface the C_{N3} coefficient (for the case $q_T^{\text{cut}} = 1$ GeV) which constitutes our best estimation. The uncertainty for each one of the C_{N3} coefficients is determined with the customary propagations of the uncertainties. The first column is used to label each particular scale choice used in figure 2.

section of less than 0.1% for the scales $\mu = M_H$ and $\mu = M_H/2$, and variations of the order of 0.3% for $\mu = M_H/4$. These variations are considerably smaller than the numerical integration error at fixed q_T^{cut} .

In table 1 and figure 2, we collect the values of C_{N3} extracted for all seven combinations of scale choices and three different values of q_T^{cut} . We note that the central value of each C_{N3} is independent of the scale (within the uncertainties), in complete agreement with eq. (2.21).




Figure 3. Total cross section of Higgs boson production, $\sigma_{N^3LO}^{H(tot.)}$, as obtained by the q_T subtraction formalism, compared with the corresponding analytical $\sigma_{N^3LO}^{H(tot.)}$ of ref. [15] (dark red dots). Green crosses with error bar denote the q_T subtraction prediction for $q_T^{cut} = 1$ GeV, red dots with error bar represents $\sigma_{N^3LO}^{H(tot.)}$ using $q_T^{cut} = 2$ GeV, and purple square dots with error bar having $q_T^{cut} = 3$ GeV. Whereas the q_T^{cut} is changed (from 1 to 3 GeV) the coefficient C_{N3} is always fixed to be the value extracted in figure 2 for $q_T^{cut} = 1$ GeV. The q_T subtraction prediction at N³LO with the C_{N3} numerical coefficient fixed to zero (using $q_T^{cut} = 1$ GeV) is shown using yellow dots with error bar. The NNLO analytical Higgs boson cross section ($\sigma_{N^3LO}^{H(tot.)}$) is represented by blue dots. All the cross sections are shown for three different scales: $\mu \equiv \mu_R = \mu_F = \{1/4, 1/2, 1\}M_H$ and horizontally displaced for better visibility. The uncertainty bars in the q_T subtraction predictions are calculated with the customary propagation of statistical uncertainties.

This scale independence of C_{N3} is unrelated to the ansatz of eq. (3.5): the terms in the right-hand side of eq. (2.21) are all scale independent and the relation between C_{N3} and $\hat{H}_g^{H;(3)}$ is defined through eqs. (2.21), (2.22) and (3.5). The uncertainties shown in figure 2 are determined using conventional error propagation and are almost entirely dominated by the size of the statistical errors of the N³LO $\sigma_H^{(fin.)}$ cross section shown in figure 1.

Since the resulting cross sections at different scale values are statistically correlated, we propose as our estimation for the C_{N3} coefficient the value obtained for $q_T^{cut} = 1$ GeV at the central scale $\mu_F = \mu_R = M_H/2$, $C_{N3} = -943 \pm 222$, which is indicated in bold typeface in table 1. The solid red central line in figure 2, and the associated red band are obtained using this single value.

The numerically extracted C_{N3} coefficient allows the total cross section to be computed at N³LO using the q_T subtraction method, which serves as a closure test of the approach and the approximations used, and allows the impact of uncertainties associated with the numerical evaluation of the ingredients to be quantified. In figure 3 we compare the fully analytical N³LO Higgs boson total cross section [15] (dark red dot) and our estimation (red dot with error bar) for three central scales, using $q_T^{cut} = 2$ GeV. The yellow dots with error bar represent our best approximation without the use of the C_{N3} coefficient (i.e.

$\sigma^{H(\text{tot.})}$ (pb)	Exact	q_T subtraction ($q_T^{\text{cut}} = 1 \text{ GeV}$)	q_T subtraction ($q_T^{\text{cut}} = 2 \text{ GeV}$)	q_T subtraction ($q_T^{\text{cut}} = 3 \text{ GeV}$)	q_T subtraction ($C_{N3} = 0$)
N ³ LO [$\mu = M_H/2$]	44.97	44.97 ± 0.21	44.98 ± 0.17	45.01 ± 0.15	45.86 ± 0.21
N ³ LO [$\mu = M_H$]	43.50	43.51 ± 0.12	43.51 ± 0.10	43.53 ± 0.09	44.08 ± 0.12
N ³ LO [$\mu = M_H/4$]	45.06	44.97 ± 0.38	44.95 ± 0.31	44.92 ± 0.28	46.44 ± 0.38
NNLO [$\mu = M_H/2$]	43.47	43.46 ± 0.02	43.46 ± 0.02	43.46 ± 0.02	43.46 ± 0.02
NNLO [$\mu = M_H$]	39.64	39.62 ± 0.02	39.62 ± 0.02	39.62 ± 0.02	39.62 ± 0.02
NNLO [$\mu = M_H/4$]	47.33	47.33 ± 0.02	47.33 ± 0.02	47.33 ± 0.02	47.33 ± 0.02

Table 2. The total cross section for Higgs boson production $\sigma^{H(\text{tot.})}$ at the LHC ($\sqrt{s} = 13 \text{ TeV}$). Results for NNLO and N³LO cross sections for three different scales $\mu = M_H/2$ (central scale), $\mu = M_H$ and $\mu = M_H/4$. The column “Exact” contains the results of ref. [15] computed with the numerical code of ref. [77] as detailed in the text. The results with the q_T subtraction method are obtained using three different values of q_T^{cut} (1, 2 and 3 GeV), and their uncertainties are calculated with the customary propagation of statistical errors. The last column shows $\sigma^{H(\text{tot.})}$ obtained with the q_T subtraction method and using $C_{N3} = 0$ at N³LO. The values of $\sigma^{H(\text{tot.})}$ reported in this table are shown in figure 3. The NNLO cross sections computed with the q_T subtraction method are obtained using $q_T^{\text{cut}} = 1 \text{ GeV}$, i.e. the variation of this parameter in the N³LO cross section is considered at N³LO-only.

$C_{N3} = 0$), that can be considered as the prediction of the q_T subtraction method in the case in which the total cross section is unknown (e.g. for Drell-Yan at N³LO). The uncertainty bars in the q_T subtraction prediction correspond to the statistical errors of the numerical computations and are mainly due to the finite contribution in eq. (3.3) at N³LO-only. The green crosses and purple squares correspond to our N³LO prediction using $q_T^{\text{cut}} = 1 \text{ GeV}$ and 3 GeV respectively. Notice that the q_T^{cut} variation is performed at N³LO-only, while the NNLO cross section is evaluated at fixed q_T^{cut} parameter. The NNLO cross section is also shown in figure 3 (blue dots) in order to put the size of the N³LO corrections in relation to the previous perturbative order. The total cross sections shown in figure 3 are reported in table 2.

4 The rapidity distribution of the Higgs boson

In this section we use the C_{N3} coefficient (extracted in section 3.2) to produce differential predictions at N³LO. In particular, we present differential results for the rapidity distribution of the Higgs boson. In section 4.1 we first estimate at NNLO the uncertainties introduced in the rapidity distribution by the procedure proposed in eq. (3.5). In section 4.2 we present the rapidity distribution at N³LO with the estimation of the uncertainties associated to the variation of the q_T^{cut} and C_{N3} parameters.

4.1 The NNLO rapidity distribution

In this section we aim to quantify the uncertainty in the approximation used in eq. (3.5). This approximation was first proposed in ref. [2] for Higgs production at NNLO. Since all the ingredients of the q_T subtraction formalism at NNLO are known in analytical form [39],




Figure 4. Comparison of the rapidity distribution between the exact result at NNLO (blue hatched) and an evaluation using the C_{N2} numerical coefficient (cross, dot and square points). We perform the comparison both at the level of (a) the full NNLO cross section and (b) for the coefficient function $\mathcal{H}^{H;(2)}$. The lower panels show the ratio to the exact result. For this particular example at NNLO, we employ the three-point scale variation: $\mu = \mu_R = \mu_F = \{M_H/4, M_H/2, M_H\}$.

it is possible to quantify the difference induced by the approximation compared to the exact result. This analysis further allows to assess the potential impact of the approximation that could be present at N³LO in section 4.2 and 4.3 below. For this quantitative study we consider the collinear functions $C_{ga}^{(1)}$ and the hard-virtual factor $H_g^{H;(1)}$ in eq. (2.20) as known. The collinear functions $C_{ga}^{(2)}$ and the first order helicity-flip functions $G_{ga}^{(1)}$ are regarded as unknown. The hard-virtual factor $H_g^{H;(2)}$ is divided in two contributions in analogy to eq. (2.22)

$$H_g^{H;(2)} \equiv \tilde{H}_g^{H;(2)} + [H_g^{H;(2)}]_{(\delta_{(1)}^{q_T})}, \quad (4.1)$$

where $[H_g^{H;(2)}]_{(\delta_{(1)}^{q_T})}$ is considered as unknown for the present NNLO study.

These so-called *unknown functions* (for this exercise) which depend on the variables z_i in eq. (2.20) are approximated with a single numerical coefficient C_{N2} proportional to $\delta(1-z_1)\delta(1-z_2)$ (the C_{N2} here was labeled as C_N in ref. [2]) in direct analogy to eq. (3.5):

$$\begin{aligned} & [H_g^{H;(2)}]_{(\delta_{(1)}^{q_T})} \delta_{ga} \delta(1-z_1) \delta_{gb} \delta(1-z_2) \\ & + C_{ga}^{(2)}(z_1) \delta_{gb} \delta(1-z_2) + \delta_{ga} \delta(1-z_1) C_{gb}^{(2)}(z_2) + G_{ga}^{(1)}(z_1) G_{gb}^{(1)}(z_2) \\ & \approx C_{N2} \delta_{ga} \delta(1-z_1) \delta_{gb} \delta(1-z_2). \end{aligned} \quad (4.2)$$

In figure 4(a) we show the rapidity distribution of the Higgs boson at NNLO computed with the exact q_T subtraction (blue hatched band) and the NNLO prediction using the

C_{N2} coefficient (dot, cross and square points). For this particular example at NNLO, we employ the three-point scale variation: $\mu = \mu_R = \mu_F = \{M_H/4, M_H/2, M_H\}$. Repeating the analysis performed for table 1 and figure 2, we obtain: $C_{N2} = 28 \pm 1$. The numerical value of the C_{N2} parameter corresponds to a specific $\tilde{H}_g^{H;(2)}$ hard coefficient:

$$\tilde{H}_g^{H;(2)} = \frac{11399}{144} + \frac{19}{8}L_t - \frac{1189}{144}N_f + \frac{2}{3}N_fL_t + \frac{83}{6}\pi^2 - \frac{5}{18}\pi^2N_f + \frac{13}{16}\pi^4 - \frac{165}{4}\zeta_3 + \frac{5}{6}N_f\zeta_3, \quad (4.3)$$

which is obtained with the same method that was used to arrive at eq. (2.23). Using this C_{N2} parameter we can produce differential predictions which are obtained *mimicking* the strategy that we intend to apply at N³LO.

In the lower panel of figure 4(a) we show the ratio to the exact NNLO result, i.e. we present the ratio for each scale. As expected, the approximation presents its best behaviour at central rapidity and the deviation from the exact results is at *per mille* level throughout the considered rapidity range of $|y_H| \leq 4$.

The study shown in figure 4(a) validates the quality of our method for the total rapidity distribution of the Higgs boson at NNLO. One could argue that a more stringent check would involve only the quantities involved in the approximation, i.e., the rapidity distribution of the second-order coefficient functions $\mathcal{H}^{H;(2)}$.

In figure 4(b) we compare the rapidity distribution for $\mathcal{H}_{\text{exact}}^{H;(2)}$ (defined in eq. (2.20)) with the approximated y_H distribution of the coefficient $\mathcal{H}_{C_{N2}}^{H;(2)}$, defined in eq. (4.2). The function $\mathcal{H}_{C_{N2}}^{H;(2)}$ approximates the exact $\mathcal{H}_{\text{exact}}^{H;(2)}$ within a precision of 2%, demonstrating the accuracy of the proposed method even at the level of individual coefficients. This directly implies that the contribution of the hard-virtual factor $H_g^{H;(2)}$ is more important than the rapidity-dependent functions $G_{ga}^{(1)}(z)$ and $C_{ga}^{(2)}(z)$ across the whole rapidity range.

We performed at NNLO variations of the q_T^{cut} value between 0.1 GeV and 3 GeV, and the NNLO cross sections (and differential distributions) present deviations within a range of size 0.26% (the largest deviation is always observed for the scale choice $\mu = M_H/4$). We consider $q_T^{\text{cut}} = 1$ GeV enough to proceed at NNLO (and as our reference value), as we can understand from table 2 at NNLO.

Summarizing, we have presented in this subsection a validation at NNLO of the approximation used at N³LO. We have performed two kinds of tests: i) a check over the observable and ii) a validation at the level of the coefficients involved in the approximation. While case ii) establishes the quality of the approach regarding the approximated particular quantities, case i) evaluates the precision of the approximation at the level of the observable which is the decisive and strongest test.

4.2 Numerical stability of the N³LO rapidity distribution

In this section, we quantify the numerical stability (as well as the involved intrinsic uncertainties) of the Higgs boson rapidity distribution at N³LO concerning the q_T^{cut} and C_{N3} parameters and the statistical uncertainties introduced by $d\sigma^{H(\text{fin.})}/dy_H$ at N³LO-only.

In figure 5 we show the rapidity distribution at N³LO obtained with the q_T subtraction method using the C_{N3} coefficient determined in section 3.2 ($C_{N3} = -943 \pm 222$). The NNLO




Figure 5. Rapidity distribution of the Higgs boson as computed using the q_T subtraction formalism at $N^3\text{LO}$. All bands include the seven-point scale variation as detailed in table 1. The red band constitutes our result with $q_T^{\text{cut}} = 2\text{ GeV}$ using the central value for the C_{N3} coefficient ($C_{N3} = -943$). The pale yellow band is obtained as the envelope between the prediction at $q_T^{\text{cut}} = 1\text{ GeV}$ and 2 GeV using $C_{N3} = -943$. The black band is computed at fixed $q_T^{\text{cut}} = 2\text{ GeV}$ taking the two extremal values of the C_{N3} coefficient according to the uncertainty ($C_{N3} = -943 \pm 222$), and performing seven-point scale variation as described in the text.

prediction is always computed with $q_T^{\text{cut}} = 1\text{ GeV}$. The red band in figure 5 shows the size of the seven-point scale variation for $q_T^{\text{cut}} = 2\text{ GeV}$.

The pale yellow band is calculated as the envelope of the scale variation bands for two different values of q_T^{cut} : 1 GeV and 2 GeV . Therefore, the pale yellow band in figure 5 can be taken as an estimate of the uncertainty due to the variation of the q_T^{cut} parameters at $N^3\text{LO}$. In figure 3 (and table 2), we observed that the total cross section (for the three central scales) is rather stable as a function of the q_T^{cut} value. The variations of the $N^3\text{LO}$ cross sections were at the *per mille* level of accuracy if we consider $q_T^{\text{cut}} = 2 \pm 1\text{ GeV}$, which is far better than the associated statistical uncertainty (see table 2). The uncertainty estimate due to the q_T^{cut} variation performed in figure 5, which is differential in the Higgs-boson rapidity, confirms the stability of the total cross section reported in table 2. The rapidity distribution is almost insensitive to the change in the q_T^{cut} parameter in the region where the bulk of the cross section is concentrated ($|y_H| \leq 3.6$). At large rapidities ($|y_H| \sim 4$), where the overall contribution to the total cross section is less than 0.5% , we found the largest deviations. Such deviations are mainly related to the numerical uncertainties from $d\sigma^{H(\text{fin.})}/dy_H$ at $N^3\text{LO}$ -only.




Figure 6. Rapidity distribution of the Higgs boson computed using the q_T subtraction formalism up to $N^3\text{LO}$. The seven-point scale variation bands (as stated in table 1) of the LO, NLO, NNLO and $N^3\text{LO}$ (C_{N3}) results are as follows: LO (pale grey fill), NLO (green fill), NNLO (blue hatched) and $N^3\text{LO}$ (C_{N3}) (red cross-hatched). The central scale ($\mu = M_H/2$) at each perturbative order (except LO) is shown with solid lines. In the lower panel, the ratio to the NNLO prediction is shown. While the bands for the predictions at LO, NLO and NNLO are computed with the seven scales as detailed in the text, the $N^3\text{LO}$ (C_{N3}) band is obtained after considering also the uncertainties due to the variation of the q_T^{cut} and the C_{N3} coefficient in the $N^3\text{LO}$ -only contribution.

Finally, we consider the uncertainty introduced by the statistical errors of the C_{N3} coefficient. The black band in figure 5 is obtained as the envelope of the seven-point scale variation at $q_T^{\text{cut}} = 2 \text{ GeV}$ now considering for each scale the two extremal C_{N3} coefficients corresponding to its maximum and minimum statistical deviations: $C_{N3} = \{-1165, -721\}$. The envelope is therefore taken from a total of 14 rapidity distributions (two extremal predictions for each one of the seven scales). The net effect of this C_{N3} variation result in an overall enlargement of the red band at $q_T^{\text{cut}} = 2 \text{ GeV}$. Our final uncertainty estimate in the rapidity distribution of the Higgs boson at $N^3\text{LO}$ is computed as the envelope of three bands: seven-point scale variation only, combined with q_T^{cut} variation, and combined with C_{N3} variation.

4.3 The rapidity distribution of the Higgs boson at $N^3\text{LO}$

In this section we present our predictions for the Higgs boson rapidity distributions at the LHC, applying the $N^3\text{LO}$ q_T subtraction method presented in section 2. The setup of the calculation is summarised in section 3.2.

Figure 6 shows the rapidity distribution of the Higgs boson at LO (pale grey fill), NLO (green fill), NNLO (blue hatched) and N³LO (red cross-hatched). The central scale ($\mu = M_H/2$) is shown as a solid line while the bands correspond to the envelope of seven-point scale variation. At N³LO, the band additionally includes the uncertainties due to q_T^{cut} and C_{N3} as described in section 4.2. Going from LO to NNLO, the scale $\mu = M_H/2$ is always at the center of the respective scale variation band in figure 6. The central prediction at N³LO, on the other hand, almost coincides with the upper edge of the band, as was already observed for the total cross section [14, 15], see table 2 and figure 3. Figures 3 and 6 respectively show a substantial reduction in the size of the scale variation band at N³LO both in the total cross section and in differential distributions.

In the central rapidity region of $|y_H| \leq 3.6$, the impact of the N³LO corrections on the NNLO result is almost independent of y_H with a flat K -factor about 1.034 for the central scale choice. The combined theoretical uncertainty at N³LO is at most of $\pm 5\%$ level with respect to the central scale choice. The uncertainty on the y_H distribution is reduced by more than a factor of 1/2 by going from NNLO to N³LO. The N³LO uncertainty band lies fully within the scale variation band at NNLO, exhibiting a stable perturbative behaviour. The only exception is the very high rapidity region, where the q_T^{cut} uncertainty becomes the dominant source for the size of the N³LO band as shown in figure 5.

The N³LO corrections to the Higgs boson rapidity distribution have been investigated in refs. [17, 18] employing a threshold expansion. The first two leading terms in the threshold expansion were computed in ref. [17], which agrees well with our calculation for the rapidity region $y_H < 0.5$ despite different choices of PDFs and scale-variation prescriptions. Both calculations display a considerable reduction of scale uncertainties going from NNLO to N³LO in this central rapidity region. For the rapidity region $y_H > 1$, however, larger differences are observed between the two calculations, where the results using the q_T subtraction formalism generally yield smaller N³LO corrections (within the NNLO scale uncertainty band). Most recently, the calculation of the threshold expansion including the first six terms was completed in ref. [18], which exhibits a stabilisation of the perturbative series together with a reduction of scale uncertainties. Comparing figure 6 with the results obtained in ref. [18], we observe very good agreement between the two calculations.

5 Conclusions and outlook

In this paper we have performed a detailed study of Higgs boson production at the LHC using the q_T subtraction formalism at N³LO. We systematically describe the q_T subtraction formalism for a generic colourless and massive system $F(\{q_i\})$ produced at hadron colliders. Fully differential cross sections for this type of final state system are separated into $\delta(q_T)$ and $q_T \neq 0$ contributions. The contribution for $q_T \neq 0$ is calculated, using a phase space cut-off q_T^{cut} , as the difference between $F(\{q_i\}) + \text{jet(s)}$ production and q_T counterterms. Specifically, we use the NNLOJET package to compute NNLO Higgs-plus-jet production and expand the Sudakov form factor in the hard resummation scheme to the matching order for the corresponding q_T counterterms. The contribution at $\delta(q_T^2)$ is further factorized into convolutions of the Sudakov form factor, the hard-virtual function, the helicity-flip

coefficient function, the hard-collinear coefficient function as well as the PDFs (section 2). The factorization guarantees that all the process-dependent contributions proportional to a form factor are included in the hard-virtual function, which depends on both initial- and final-state particles. All other factorized contributions only depend on the initial states. Some of the factorized ingredients contributing at $\delta(q_T^2)$ are not known analytically at N³LO for the moment. We collect all analytically available contributions and approximate the unknown pieces by a constant coefficient C_{N3} which is scale- and process-independent (section 3). Using the available inclusive total cross section for N³LO Higgs production and the known pieces from the q_T subtraction formalism, we numerically extract the value of C_{N3} . By comparing the numerical values for C_{N3} using different scales and q_T^{cut} setups in the extraction, we conclude from mutually consistent results that C_{N3} is independent of the scale choice with a value obtained for $\mu = M_H/2$ and $q_T^{\text{cut}} = 1 \text{ GeV}$ of $C_{N3} = -943 \pm 222$ (section 3.2).

As a proof-of-concept implementation of the q_T subtraction method at N³LO, we calculate the total cross section and rapidity distributions for Higgs boson production at LHC using a new Monte Carlo generator HN3LO [76]. Using the extracted value of C_{N3} , we perform a closure test for the inclusive total cross section for three different scale choices and find excellent agreement with the exact results (from `ihixs 2` [77]) at the 0.2% level. For the differential rapidity distribution of the Higgs boson, we first study the systematic error from the C_{N3} approximation by considering the NNLO calculation and introducing an approximate C_{N2} . The NNLO y_H distribution exhibit per-mille level agreement between the C_{N2} approximation and the exact result, supporting the reliability of the procedure. We calculate the y_H distribution at N³LO employing a seven-point scale variation and carefully assess systematic errors arising from different q_T^{cut} and C_{N3} values. Compared to the NNLO y_H distributions, we observe a large reduction of theory uncertainties by more than 50% at N³LO. The scale variation band at N³LO stays within the NNLO band with a flat K -factor of about 1.034 in the central rapidity region ($|y_H| \leq 3.6$). Both the systematic error analysis and the phenomenological predictions confirm that our calculations at N³LO using q_T subtraction formalism are well under control. The approximation related to the C_{N3} coefficient in our approach can be easily replaced by the full analytical results once available.

With the upcoming larger data set and more accurate measurements of Higgs properties at the LHC, we prepare precise theoretical tools that could match the frontier accuracy of experimental results. More differential properties at N³LO involving the Higgs boson and its decay products can be studied using the same framework established in this paper. The current N³LO calculation, using the approximation of large top quark mass, attains a level of precision that several other contributions will need to be taken into account for a full study of precision phenomenology [78]: finite top quark mass effects, heavy-light quark interference contributions and electroweak corrections.

Acknowledgments

LC would like to thank Stefano Catani for very useful and valuable discussions. XC would like to thank Javier Mazzitelli and Hua Xing Zhu for inspiring discussions. We thank the

University of Zurich S3IT and CSCS Lugano for providing the computational resources for this project. This research was supported in part by the U.K. Science and Technology Facilities Council, by the Swiss National Science Foundation (SNF) under contracts 200020-175595 and CRSII2-160814, by the Swiss National Supercomputing Centre (CSCS) under project ID UZH10, by the Research Executive Agency (REA) of the European Union under the ERC Advanced Grant MC@NNLO (340983).

A Fixed-order expressions

The precise identification of the Sudakov form factor S_c , the hard-virtual function $H_g^{F=H}$ and the hard-collinear coefficient functions, C_{ga} and G_{ga} is not unique, and the resummation formula (2.10) is invariant under “resummation scheme” transformations [32]:

$$\begin{aligned} H_c^F(\alpha_s) &\rightarrow H_c^F(\alpha_s) [h(\alpha_s)]^{-1}, \\ B_c(\alpha_s) &\rightarrow B_c(\alpha_s) - \beta(\alpha_s) \frac{d \ln h(\alpha_s)}{d \ln \alpha_s}, \\ C_{ab}(\alpha_s, z) &\rightarrow C_{ab}(\alpha_s, z) [h(\alpha_s)]^{1/2}, \\ G_{ab}(\alpha_s, z) &\rightarrow G_{ab}(\alpha_s, z) [h(\alpha_s)]^{1/2}. \end{aligned} \quad (\text{A.1})$$

This invariance can easily be proven by using the following renormalization-group identity:

$$h(\alpha_s(b_0^2/b^2)) = h(\alpha_s(M^2)) \exp \left\{ - \int_{b_0^2/b^2}^{M^2} \frac{dq^2}{q^2} \beta(\alpha_s(q^2)) \frac{d \ln h(\alpha_s(q^2))}{d \ln \alpha_s(q^2)} \right\}, \quad (\text{A.2})$$

which is valid for any perturbative function $h(\alpha_s)$. Notice that eq. (A.2) establishes the evolution of the perturbative functions from the scale $q^2 = b_0^2/b^2$ to $q^2 = M^2$. The QCD β -function and its corresponding n -th order coefficient β_n are defined as

$$\frac{d \ln \alpha_s(\mu^2)}{d \ln \mu^2} = \beta(\alpha_s(\mu^2)) = - \sum_{n=0}^{+\infty} \beta_n \left(\frac{\alpha_s}{\pi} \right)^{n+1}. \quad (\text{A.3})$$

The explicit expression of the first three coefficients [36, 37], β_0 , β_1 and β_2 read

$$\begin{aligned} \beta_0 &= \frac{1}{12} (11C_A - 2N_f), \quad \beta_1 = \frac{1}{24} (17C_A^2 - 5C_A N_f - 3C_F N_f), \\ \beta_2 &= \frac{1}{64} \left(\frac{2857}{54} C_A^3 - \frac{1415}{54} C_A^2 N_f - \frac{205}{18} C_A C_F N_f + C_F^2 N_f + \frac{79}{54} C_A N_f^2 + \frac{11}{9} C_F N_f^2 \right), \end{aligned} \quad (\text{A.4})$$

where N_f is the number of massless QCD flavours and the $SU(N_c)$ colour factors are $C_A = N_c$ and $C_F = (N_c^2 - 1)/(2N_c)$.

Throughout this paper we always use the *hard resummation scheme* [35] to report explicit expressions for the perturbative expansion of these individual coefficients. The *hard resummation scheme* states that all the contributions proportional to $\delta(1 - z)$ are associated with the hard-virtual functions H_c^F . This directly implies that H_c^F is process dependent. The collinear C_{ab} and G_{ab} functions and the resummation coefficients A_c and B_c are independent of the final state system F .

The truncation of eq. (2.10) at a given fixed order requires the explicit knowledge of resummation coefficients and hard collinear coefficient functions. For $F = H$ at NLO, the knowledge of the coefficients $A_g^{(1)}$, $B_g^{(1)}$, $C_{ga}^{(1)}$ ($a = q, \bar{q}, g$) and $H_g^{H;(1)}$ are sufficient to compute the inclusive total cross section and differential distributions. Assuming that the Higgs boson couples to a single heavy quark of mass m_Q , the first-order coefficient $H_g^{H;(1)}$ in the hard resummation scheme is [35]

$$H_g^{H;(1)} = C_A \pi^2 / 2 + c_H(m_Q). \quad (\text{A.5})$$

The function $c_H(m_Q)$, which depends on the NLO virtual corrections of the Born subprocess, is given in eq. (B.2) of ref. [54]. In the limit $m_Q \rightarrow \infty$, the function c_H becomes

$$c_H(m_Q) \longrightarrow \frac{5C_A - 3C_F}{2}. \quad (\text{A.6})$$

Therefore, the complete set of coefficients necessary to compute Higgs boson production (in the limit in which the mass of the top quark $Q = t$ is larger than any other scale involved in the process) at NLO are

$$\begin{aligned} A_g^{(1)} &= C_A, & B_g^{(1)} &= -\frac{1}{6}(11C_A - 2N_f), & H_g^{H;(1)} &= \frac{1}{2}(C_A(\pi^2 + 5) - 3C_F), \\ C_{gg}^{(1)}(z) &= 0, & C_{ga}^{(1)}(z) &= \frac{1}{2}C_F z & [a = q, \bar{q}]. \end{aligned} \quad (\text{A.7})$$

The coefficients $A_g^{(1)}$ and $B_g^{(1)}$ are process *and* resummation scheme independent. The collinear functions $C_{ga}^{(1)}$ ($a = q, \bar{q}, g$) are process independent, while $H_g^{H;(1)}$ depends on the final-state system ($F = H$). Together, they depend on the resummation scheme in such a way to ensure the resummation scheme independence of eq. (2.10) at NLO. In ref. [38] was shown that the NLO hard-virtual coefficient $H_c^{F;(1)}$ is explicitly related to $d\hat{\sigma}_{\text{LO}}^F$ and to the IR finite part of the NLO virtual correction to the Born cross section.

At NNLO, the coefficients $A_g^{(2)}$ and $B_g^{(2)}$ are needed [2, 35, 38],

$$A_g^{(2)} = \frac{1}{2} C_A \left[\left(\frac{67}{18} - \frac{\pi^2}{6} \right) C_A - \frac{5}{9} N_f \right], \quad B_g^{(2)} = \frac{\gamma_g^{(1)}}{16} + \beta_0 C_A \zeta_2, \quad (\text{A.8})$$

where $\gamma_g^{(1)}$ is the coefficient of the $\delta(1-z)$ term in the NLO gluon splitting function [44, 45], which reads

$$\gamma_g^{(1)} = \left(-\frac{64}{3} - 24\zeta_3 \right) C_A^2 + \frac{16}{3} C_A N_f + 4 C_F N_f. \quad (\text{A.9})$$

The coefficient $A_g^{(2)}$ does not depend on the resummation scheme whereas $B_g^{(2)}$ in eq. (A.8) is valid in the hard resummation scheme and both coefficients are process independent.

The general structure of the hard-virtual coefficients H_c^F has been established in ref. [35]. Although H_c^F is in principle process dependent, ref. [35] showed it can be directly related in a universal way to the IR finite part of the all-order virtual amplitude of the corresponding partonic subprocess $c\bar{c} \rightarrow F$. The relationship between H_c^F and the all-order virtual correction to the partonic subprocess $c\bar{c} \rightarrow F$ has been made explicit up

to NNLO and is based on the definition of universal subtraction operators that cancel the IR divergences of the two-loop (NNLO) virtual corrections to the Born cross section [47]. These universal second-order operators contain an IR finite term of soft origin ($\delta_{qT}^{(1)}$) that only depends on the initial-state partons [35].

In the case of Higgs boson production, the hard-virtual factor $H_g^{F=H;(2)}$ in the large- m_t limit (in the hard resummation scheme) is given by [39]

$$\begin{aligned} H_g^{H;(2)} = & C_A^2 \left(\frac{3187}{288} + \frac{7}{8} L_t + \frac{157}{72} \pi^2 + \frac{13}{144} \pi^4 - \frac{55}{18} \zeta_3 \right) + C_A C_F \left(-\frac{145}{24} - \frac{11}{8} L_t - \frac{3}{4} \pi^2 \right) \\ & + \frac{9}{4} C_F^2 - \frac{5}{96} C_A - \frac{1}{12} C_F - C_A N_f \left(\frac{287}{144} + \frac{5}{36} \pi^2 + \frac{4}{9} \zeta_3 \right) \\ & + C_F N_f \left(-\frac{41}{24} + \frac{1}{2} L_t + \zeta_3 \right), \end{aligned} \quad (\text{A.10})$$

where $L_t = \ln(M^2/m_t^2)$. The two-loop scattering amplitude [46] used in the computation of $H_g^{F=H;(2)}$ includes corrections to the large- m_t approximation.

Due to the large size of the expressions for $C_{ab}^{(2)}(z)$, we refrain from explicitly quoting them here and instead refer to eqs. (37)–(40) of ref. [35] using the full results of refs. [39, 40]. These collinear coefficients $C_{ab}^{(2)}$ have been independently computed in refs. [41–43].

At NNLO, in eq. (2.13) the first order $G_{ga}^{(1)}$ helicity-flip functions are required which read [34]

$$G_{ga}^{(1)}(z) = C_a \frac{1-z}{z} \quad a = q, \bar{q}, g, \quad (\text{A.11})$$

where $C_{q;\bar{q}} = C_F$ and $C_g = C_A$. The first-order functions $G_{ga}^{(1)}$ are resummation-scheme independent and do not depend on the final-state system F .

At N³LO, the numerical implementation of eq. (2.10) requires the following ingredients: $A_g^{(3)}$, $B_g^{(3)}$, $C_{ga}^{(3)}$, $G_{ga}^{(2)}$ ($a = q, \bar{q}, g$) and $H_g^{H;(3)}$. The coefficient $A_g^{(3)}$ [58] reads

$$\begin{aligned} A_g^{(3)} = & C_A^3 \left(\frac{245}{96} - \frac{67}{36} \zeta_2 + \frac{11}{24} \zeta_3 + \frac{11}{20} \zeta_2^2 \right) + C_A C_F N_f \left(-\frac{55}{96} + \frac{1}{2} \zeta_3 \right) - C_A N_f^2 \frac{1}{108} \\ & + C_A^2 N_f \left(-\frac{209}{432} + \frac{5}{18} \zeta_2 - \frac{7}{12} \zeta_3 \right) + \beta_0 C_A^2 \left(\frac{101}{27} - \frac{7}{2} \zeta_3 \right) - \beta_0 C_A N_f \frac{14}{27}. \end{aligned} \quad (\text{A.12})$$

The explicit expression of the $B_c^{(3)}$ ($a = q, g$) coefficients in the hard scheme can be computed from refs. [55, 56]. In the particular case of the gluon channel then in the hard resummation scheme, we obtain

$$\begin{aligned} B_g^{(3)} = & -\frac{2133}{64} + \frac{3029}{576} N_f - \frac{349}{1728} N_f^2 + \frac{109}{6} \pi^2 - \frac{283}{144} \pi^2 N_f + \frac{5}{108} \pi^2 N_f^2 - \frac{253}{160} \pi^4 \\ & + \frac{23}{240} \pi^4 N_f - \frac{843}{8} \zeta_3 + 2 \zeta_3 N_f + \frac{1}{6} \zeta_3 N_f^2 + \frac{9}{4} \pi^2 \zeta_3 + \frac{135}{2} \zeta_5. \end{aligned} \quad (\text{A.13})$$

B Convolutions at N³LO

The numerical implementation of eq. (2.10) requires the computation of several convolutions between splitting functions, collinear and helicity-flip functions. In principle, taking

(i)	$\gamma_{ga}^{(1)} \otimes \gamma_{ab}^{(1)} \otimes \gamma_{bg}^{(1)}$	(ii)	$\gamma_{ga}^{(1)} \otimes \gamma_{ab}^{(1)} \otimes \gamma_{bq}^{(1)}$
(iii)	$\gamma_{ga}^{(1)} \otimes \gamma_{ag}^{(2)}$	(iv)	$\gamma_{ga}^{(1)} \otimes \gamma_{aq}^{(2)}$
(v)	$\gamma_{ga}^{(2)} \otimes \gamma_{ag}^{(1)}$	(vi)	$\gamma_{ga}^{(2)} \otimes \gamma_{aq}^{(1)}$
(vii)	$C_{ga}^{(1)} \otimes \gamma_{ag}^{(2)}$	(viii)	$C_{ga}^{(1)} \otimes \gamma_{aq}^{(2)}$
(ix)	$C_{ga}^{(2)} \otimes \gamma_{ag}^{(1)}$	(x)	$C_{ga}^{(2)} \otimes \gamma_{aq}^{(1)}$
(xi)	$G_{ga}^{(1)} \otimes \gamma_{ag}^{(1)}$	(xii)	$G_{ga}^{(1)} \otimes \gamma_{aq}^{(1)}$

Table 3. Convolutions appearing at the N³LO-only between the collinear $C_{ab}^{(n)}$, the helicity-flip $G_{ab}^{(n)}$ and the splitting functions $\gamma_{ab}^{(n)}$ ($n = 1, 2$). The repeated subindices a and b imply a sum over the parton flavors q, \bar{q}, g . The first and last subindices denote the partonic channel in which they are contributing, i.e. the convolutions in the first column are used in the gg partonic channel whereas the second (and last) column is for the qg and gq partonic channels.

the N -moments of the functions involved in the calculation, one can avoid the use of convolutions, since in N -space they correspond to simple products. However, the numerical implementation of eq. (2.10) in the Monte Carlo code **HN3LO** was carried out in the z -space (e.g. as in the codes **HNNLO** [1], **DYNNLO** [79], **2 γ NNLO** [80], etc.), and therefore the new third order convolutions have to be calculated as well.

The convolutions in eqs. (2.20), (2.21), (3.5) and (4.2) between two functions ($f(z)$ and $g(z)$) of the variable z are defined through the following integral

$$(f \otimes g)(z) \equiv \int_z^1 \frac{dy}{y} f\left(\frac{z}{y}\right) g(y). \quad (\text{B.1})$$

In the case of processes initiated by gluon fusion, the complete list of third order convolutions to be calculated can be found in table 3. All the remaining convolutions in eq. (2.10) at N³LO already contributed to the previous orders and they are regarded as known.

The symbol $\gamma_{ab}^{(n)}$ in table 3 denotes the usual splitting functions of n -th order and they contribute to eq. (2.10) since the PDFs have to be evolved from the scale b_0^2/b^2 to the factorization scale μ_F . The first three rows in eq. (3) were calculated in ref. [82] and cross-checked with a dedicated computation for the results presented in this paper. The public **Mathematica** package **MT** [81] is used to calculate the necessary convolutions (i)–(vi) in ref. [82], which can be further expressed in terms of *harmonic polylogarithms* (HPLs) [85] using the **Mathematica** package **HPL** [84]. The remaining convolutions in eqs. (vii)–(xii) of table 3 were computed for this work. The **MT** [81] package is not able to solve all the convolutions of weight 3 and 4 that are needed in (vii)–(xii). For instance, the **MT** package cannot handle convolutions in which their result has to be expressed in terms of multiple polylogarithms (or *Goncharov polylogarithms* GPLs) [83, 86, 87] as it is the case when the collinear functions $C_{gj}^{(2)}$ are involved. For those, we have computed the convolutions (vii)–(xii) with a newly developed code **Convo**, which is able to provide results in terms of GPLs and also can handle terms that are individually divergent, but finite after addition.

(a)	$G(\frac{z}{1+z}, 0, 0, 1; \frac{1}{2})$	(b)	$G(1, 0, 0, -z; z)$	(c)	$G(0, 1, 0, -1; z)$
(d)	$G(0, 1, 0, z; 1)$	(e)	$G(0, 1, z, 0; 1)$	(f)	$G(0, z, 1, 0; 1)$
(g)	$G(-z, 0, z, 0; 1)$	(h)	$G(0, 1, 0, -z; z)$	(i)	$G(0, 1, -z, -z; z)$
(j)	$G(-z, 1, 0, 0; 1)$	(k)	$G(-z, 1, 0, 0; z)$	(l)	$G(-z, 0, 0, z; 1)$

Table 4. Basis for the GPLs used in the numerical implementation of the convolutions listed in table 3.

The multiple polylogarithms can be defined recursively, for $n \geq 0$, via the iterated integral [83, 86, 87]

$$G(a_1, \dots, a_n; z) = \int_0^z \frac{dt}{t - a_1} G(a_2, \dots, a_n; t), \quad (\text{B.2})$$

with $G(z) = G(; z) = 1$ (an exception being when $z = 0$ in which case we put $G(0) = 0$) and with $a_i \in \mathbb{C}$ are chosen constants and z is a complex variable. For the convolutions in table 3 the variable z and the weights a_1, \dots, a_n are all real constants.

From the convolutions in table 3 we quote some examples which appear as building blocks in the computation of eqs. (vii)–(xii),

$$\left\{ D_0[1 - y]; \frac{1}{y}; 1; y; y^2 \right\} \otimes \left(\frac{f(y)}{1 + y} \right), \quad (\text{B.3})$$

with

$$f(y) = \left\{ \text{Li}_3 \left(\frac{1}{1 + y} \right); \text{Li}_3(\pm y); \text{Li}_2(\pm y); \text{Li}_2(1 - y); \text{Li}_2(\pm y) \ln(y); \right. \\ \left. \ln^2(1 + y) \ln(y); \ln(1 + y) \ln^2(y) \right\}, \quad (\text{B.4})$$

where the *plus* distribution $D_0[1 - z]$ is defined as usual

$$\int_0^1 dz f(z) D_0[1 - z] = \int_0^1 dz \frac{f(z)}{(1 - z)_+} = \int_0^1 \frac{dz}{1 - z} (f(z) - f(1)). \quad (\text{B.5})$$

After performing all the convolutions listed in table 3, their final expressions (each one of the convolutions) are finite in the domain $z \in (0, 1)$. Even more, convolutions evaluated in the domain $z \in (0, 1)$ produce results in \mathbb{R} . It is possible to write the expressions in table 3 (after simplifying) in terms of twelve GPLs that are not reducible to polylogarithmic functions of type $\text{Li}_n(z)$, and cannot be combined (e.g. through the *shuffle* algebra) with other GPLs in order to produce simpler results. The list of the irreducible GPLs is presented in table 4. All remaining GPLs appearing in the convolutions of table 3 can be related to the set given in table 4 using the results of refs. [84, 88, 89] and performing the customary *shuffle* algebra. The numerical implementation of the GPLs in table 4 was made using the package `GiNaC` [90, 91]. The basis of GPLs in table 4 is not unique, but sufficient for numerical evaluation.

An example of a third order convolution is the following integral

$$\begin{aligned}
 \left(\frac{\text{Li}_3(y)}{1+y} \otimes \text{D}_0[1-y] \right) (z) &= \int_z^1 \frac{dy}{y+z} \text{Li}_3\left(\frac{z}{y}\right) \frac{1}{(1-y)_+} \\
 &= \frac{1}{1+z} \left(-\zeta_3 G(0; z) + \frac{i\pi^3}{6} G(0; z) + \frac{\pi^2}{3} G(-z; 1) G(0; z) - i\pi G(-z, 0; 1) G(0; z) \right. \\
 &\quad - G(-z, 0, 0; 1) G(0; z) + \frac{i\pi\zeta_3}{4} + \frac{\pi^2}{3} G(0, 1; z) + i\pi G(-z; 1) G(0, 0; z) - \frac{\pi^2}{6} G(0, 0; z) \\
 &\quad - G(-z; 1) G(0, 0, 0; z) + i\pi G(0, 0, 1; z) + G(0, 0; z) G(-z, 0; 1) - \frac{\pi^2}{3} G(-z, 0; 1) \\
 &\quad + i\pi G(-z, 0, 0; 1) - G(0, 0, 0, 1; z) - G(0, 0, 1, z; 1) - G(0, 0, z, 1; 1) - G(0, 1, 0, z; 1) \\
 &\quad \left. - G(1, 0, 0, z; z) + G(-z, 0, 0, 0; 1) - G(-z, 0, 0, z; 1) + G(-z, 0, 0, z; z) + \frac{19\pi^4}{720} \right).
 \end{aligned} \tag{B.6}$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] S. Catani and M. Grazzini, *An NNLO subtraction formalism in hadron collisions and its application to Higgs boson production at the LHC*, *Phys. Rev. Lett.* **98** (2007) 222002 [[hep-ph/0703012](#)] [[INSPIRE](#)].
- [2] G. Bozzi, S. Catani, D. de Florian and M. Grazzini, *Transverse-momentum resummation and the spectrum of the Higgs boson at the LHC*, *Nucl. Phys. B* **737** (2006) 73 [[hep-ph/0508068](#)] [[INSPIRE](#)].
- [3] R. Bonciani, S. Catani, M. Grazzini, H. Sargsyan and A. Torre, *The q_T subtraction method for top quark production at hadron colliders*, *Eur. Phys. J. C* **75** (2015) 581 [[arXiv:1508.03585](#)] [[INSPIRE](#)].
- [4] R. Boughezal, X. Liu and F. Petriello, *N -jettiness soft function at next-to-next-to-leading order*, *Phys. Rev. D* **91** (2015) 094035 [[arXiv:1504.02540](#)] [[INSPIRE](#)].
- [5] J. Gaunt, M. Stahlhofen, F.J. Tackmann and J.R. Walsh, *N -jettiness Subtractions for NNLO QCD Calculations*, *JHEP* **09** (2015) 058 [[arXiv:1505.04794](#)] [[INSPIRE](#)].
- [6] M. Czakon, *Double-real radiation in hadronic top quark pair production as a proof of a certain concept*, *Nucl. Phys. B* **849** (2011) 250 [[arXiv:1101.0642](#)] [[INSPIRE](#)].
- [7] R. Boughezal, K. Melnikov and F. Petriello, *A subtraction scheme for NNLO computations*, *Phys. Rev. D* **85** (2012) 034025 [[arXiv:1111.7041](#)] [[INSPIRE](#)].
- [8] M. Cacciari, F.A. Dreyer, A. Karlberg, G.P. Salam and G. Zanderighi, *Fully Differential Vector-Boson-Fusion Higgs Production at Next-to-Next-to-Leading Order*, *Phys. Rev. Lett.* **115** (2015) 082002 [Erratum *ibid.* **120** (2018) 139901] [[arXiv:1506.02660](#)] [[INSPIRE](#)].
- [9] A. Gehrmann-De Ridder, T. Gehrmann and E.W.N. Glover, *Antenna subtraction at NNLO*, *JHEP* **09** (2005) 056 [[hep-ph/0505111](#)] [[INSPIRE](#)].
- [10] A. Daleo, T. Gehrmann and D. Maître, *Antenna subtraction with hadronic initial states*, *JHEP* **04** (2007) 016 [[hep-ph/0612257](#)] [[INSPIRE](#)].

- [11] J. Currie, E.W.N. Glover and S. Wells, *Infrared Structure at NNLO Using Antenna Subtraction*, *JHEP* **04** (2013) 066 [[arXiv:1301.4693](#)] [[INSPIRE](#)].
- [12] K.G. Chetyrkin, J.H. Kuhn and A. Kwiatkowski, *QCD corrections to the e^+e^- cross-section and the Z boson decay rate*, [hep-ph/9503396](#) [[INSPIRE](#)].
- [13] J.A.M. Vermaseren, A. Vogt and S. Moch, *The third-order QCD corrections to deep-inelastic scattering by photon exchange*, *Nucl. Phys. B* **724** (2005) 3 [[hep-ph/0504242](#)] [[INSPIRE](#)].
- [14] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog and B. Mistlberger, *Higgs Boson Gluon-Fusion Production in QCD at Three Loops*, *Phys. Rev. Lett.* **114** (2015) 212001 [[arXiv:1503.06056](#)] [[INSPIRE](#)].
- [15] B. Mistlberger, *Higgs boson production at hadron colliders at N^3 LO in QCD*, *JHEP* **05** (2018) 028 [[arXiv:1802.00833](#)] [[INSPIRE](#)].
- [16] F.A. Dreyer and A. Karlberg, *Vector-Boson Fusion Higgs Production at Three Loops in QCD*, *Phys. Rev. Lett.* **117** (2016) 072001 [[arXiv:1606.00840](#)] [[INSPIRE](#)].
- [17] F. Dulat, B. Mistlberger and A. Pelloni, *Differential Higgs production at N^3 LO beyond threshold*, *JHEP* **01** (2018) 145 [[arXiv:1710.03016](#)] [[INSPIRE](#)].
- [18] F. Dulat, B. Mistlberger and A. Pelloni, *Precision Predictions at N^3 LO for the Higgs Boson Rapidity Distribution at the LHC*, [arXiv:1810.09462](#) [[INSPIRE](#)].
- [19] J. Currie, T. Gehrmann, E.W.N. Glover, A. Huss, J. Niehues and A. Vogt, *N^3 LO corrections to jet production in deep inelastic scattering using the Projection-to-Born method*, *JHEP* **05** (2018) 209 [[arXiv:1803.09973](#)] [[INSPIRE](#)].
- [20] Y.L. Dokshitzer, D. Diakonov and S.I. Troian, *On the Transverse Momentum Distribution of Massive Lepton Pairs*, *Phys. Lett.* **79B** (1978) 269 [[INSPIRE](#)].
- [21] Y.L. Dokshitzer, D. Diakonov and S.I. Troian, *Hard Processes in Quantum Chromodynamics*, *Phys. Rept.* **58** (1980) 269 [[INSPIRE](#)].
- [22] G. Parisi and R. Petronzio, *Small Transverse Momentum Distributions in Hard Processes*, *Nucl. Phys. B* **154** (1979) 427 [[INSPIRE](#)].
- [23] G. Curci, M. Greco and Y. Srivastava, *QCD Jets From Coherent States*, *Nucl. Phys. B* **159** (1979) 451 [[INSPIRE](#)].
- [24] J.C. Collins and D.E. Soper, *Back-To-Back Jets in QCD*, *Nucl. Phys. B* **193** (1981) 381 [*Erratum* *ibid.* **B 213** (1983) 545] [[INSPIRE](#)].
- [25] J.C. Collins and D.E. Soper, *Back-To-Back Jets: Fourier Transform from B to K-Transverse*, *Nucl. Phys. B* **197** (1982) 446 [[INSPIRE](#)].
- [26] J. Kodaira and L. Trentadue, *Summing Soft Emission in QCD*, *Phys. Lett.* **112B** (1982) 66 [[INSPIRE](#)].
- [27] J. Kodaira and L. Trentadue, *Soft Gluon Effects In Perturbative Quantum Chromodynamics*, SLAC-PUB-2934 [[INSPIRE](#)].
- [28] J. Kodaira and L. Trentadue, *Single Logarithm Effects in electron-Positron Annihilation*, *Phys. Lett.* **123B** (1983) 335 [[INSPIRE](#)].
- [29] J.C. Collins, D.E. Soper and G.F. Sterman, *Transverse Momentum Distribution in Drell-Yan Pair and W and Z Boson Production*, *Nucl. Phys. B* **250** (1985) 199 [[INSPIRE](#)].

- [30] S. Catani, E. D’Emilio and L. Trentadue, *The Gluon Form-factor to Higher Orders: Gluon Gluon Annihilation at Small Q^- transverse*, *Phys. Lett. B* **211** (1988) 335 [INSPIRE].
- [31] D. de Florian and M. Grazzini, *Next-to-next-to-leading logarithmic corrections at small transverse momentum in hadronic collisions*, *Phys. Rev. Lett.* **85** (2000) 4678 [[hep-ph/0008152](#)] [INSPIRE].
- [32] S. Catani, D. de Florian and M. Grazzini, *Universality of nonleading logarithmic contributions in transverse momentum distributions*, *Nucl. Phys. B* **596** (2001) 299 [[hep-ph/0008184](#)] [INSPIRE].
- [33] W. Bizoń, P.F. Monni, E. Re, L. Rottoli and P. Torrielli, *Momentum-space resummation for transverse observables and the Higgs p_\perp at $N^3LL+NNLO$* , *JHEP* **02** (2018) 108 [[arXiv:1705.09127](#)] [INSPIRE].
- [34] S. Catani and M. Grazzini, *QCD transverse-momentum resummation in gluon fusion processes*, *Nucl. Phys. B* **845** (2011) 297 [[arXiv:1011.3918](#)] [INSPIRE].
- [35] S. Catani, L. Cieri, D. de Florian, G. Ferrera and M. Grazzini, *Universality of transverse-momentum resummation and hard factors at the NNLO*, *Nucl. Phys. B* **881** (2014) 414 [[arXiv:1311.1654](#)] [INSPIRE].
- [36] O.V. Tarasov, A.A. Vladimirov and A. Yu. Zharkov, *The Gell-Mann-Low Function of QCD in the Three Loop Approximation*, *Phys. Lett.* **93B** (1980) 429 [INSPIRE].
- [37] S.A. Larin and J.A.M. Vermaseren, *The three loop QCD β -function and anomalous dimensions*, *Phys. Lett. B* **303** (1993) 334 [[hep-ph/9302208](#)] [INSPIRE].
- [38] D. de Florian and M. Grazzini, *The structure of large logarithmic corrections at small transverse momentum in hadronic collisions*, *Nucl. Phys. B* **616** (2001) 247 [[hep-ph/0108273](#)] [INSPIRE].
- [39] S. Catani and M. Grazzini, *Higgs Boson Production at Hadron Colliders: Hard-Collinear Coefficients at the NNLO*, *Eur. Phys. J. C* **72** (2012) 2013 [Erratum *ibid.* **C 72** (2012) 2132] [[arXiv:1106.4652](#)] [INSPIRE].
- [40] S. Catani, L. Cieri, D. de Florian, G. Ferrera and M. Grazzini, *Vector boson production at hadron colliders: hard-collinear coefficients at the NNLO*, *Eur. Phys. J. C* **72** (2012) 2195 [[arXiv:1209.0158](#)] [INSPIRE].
- [41] T. Gehrmann, T. Luebbert and L.L. Yang, *Transverse parton distribution functions at next-to-next-to-leading order: the quark-to-quark case*, *Phys. Rev. Lett.* **109** (2012) 242003 [[arXiv:1209.0682](#)] [INSPIRE].
- [42] T. Gehrmann, T. Luebbert and L.L. Yang, *Calculation of the transverse parton distribution functions at next-to-next-to-leading order*, *JHEP* **06** (2014) 155 [[arXiv:1403.6451](#)] [INSPIRE].
- [43] M.G. Echevarria, I. Scimemi and A. Vladimirov, *Unpolarized Transverse Momentum Dependent Parton Distribution and Fragmentation Functions at next-to-next-to-leading order*, *JHEP* **09** (2016) 004 [[arXiv:1604.07869](#)] [INSPIRE].
- [44] G. Curci, W. Furmanski and R. Petronzio, *Evolution of Parton Densities Beyond Leading Order: The Nonsinglet Case*, *Nucl. Phys. B* **175** (1980) 27 [INSPIRE].
- [45] W. Furmanski and R. Petronzio, *Singlet Parton Densities Beyond Leading Order*, *Phys. Lett.* **97B** (1980) 437 [INSPIRE].

- [46] R.V. Harlander and K.J. Ozeren, *Top mass effects in Higgs production at next-to-next-to-leading order QCD: Virtual corrections*, *Phys. Lett. B* **679** (2009) 467 [[arXiv:0907.2997](#)] [[INSPIRE](#)].
- [47] S. Catani, *The singular behavior of QCD amplitudes at two loop order*, *Phys. Lett. B* **427** (1998) 161 [[hep-ph/9802439](#)] [[INSPIRE](#)].
- [48] R.P. Kauffman, *Higher order corrections to Higgs boson p_T* , *Phys. Rev. D* **45** (1992) 1512 [[INSPIRE](#)].
- [49] D. de Florian, G. Ferrera, M. Grazzini and D. Tommasini, *Transverse-momentum resummation: Higgs boson production at the Tevatron and the LHC*, *JHEP* **11** (2011) 064 [[arXiv:1109.2109](#)] [[INSPIRE](#)].
- [50] T. Becher, M. Neubert and D. Wilhelm, *Higgs-Boson Production at Small Transverse Momentum*, *JHEP* **05** (2013) 110 [[arXiv:1212.2621](#)] [[INSPIRE](#)].
- [51] D. Neill, I.Z. Rothstein and V. Vaidya, *The Higgs Transverse Momentum Distribution at NNLL and its Theoretical Errors*, *JHEP* **12** (2015) 097 [[arXiv:1503.00005](#)] [[INSPIRE](#)].
- [52] X. Chen et al., *Precise QCD Description of the Higgs Boson Transverse Momentum Spectrum*, *Phys. Lett. B* **788** (2019) 425 [[arXiv:1805.00736](#)] [[INSPIRE](#)].
- [53] W. Bizoń et al., *Fiducial distributions in Higgs and Drell-Yan production at $N^3LL+NNLO$* , *JHEP* **12** (2018) 132 [[arXiv:1805.05916](#)] [[INSPIRE](#)].
- [54] M. Spira, A. Djouadi, D. Graudenz and P.M. Zerwas, *Higgs boson production at the LHC*, *Nucl. Phys. B* **453** (1995) 17 [[hep-ph/9504378](#)] [[INSPIRE](#)].
- [55] Y. Li and H.X. Zhu, *Bootstrapping Rapidity Anomalous Dimensions for Transverse-Momentum Resummation*, *Phys. Rev. Lett.* **118** (2017) 022004 [[arXiv:1604.01404](#)] [[INSPIRE](#)].
- [56] A.A. Vladimirov, *Correspondence between Soft and Rapidity Anomalous Dimensions*, *Phys. Rev. Lett.* **118** (2017) 062001 [[arXiv:1610.05791](#)] [[INSPIRE](#)].
- [57] S. Catani, L. Cieri, D. de Florian, G. Ferrera and M. Grazzini, *Threshold resummation at N^3LL accuracy and soft-virtual cross sections at N^3LO* , *Nucl. Phys. B* **888** (2014) 75 [[arXiv:1405.4827](#)] [[INSPIRE](#)].
- [58] T. Becher and M. Neubert, *Drell-Yan Production at Small q_T , Transverse Parton Distributions and the Collinear Anomaly*, *Eur. Phys. J. C* **71** (2011) 1665 [[arXiv:1007.4005](#)] [[INSPIRE](#)].
- [59] F. Wilczek, *Decays of Heavy Vector Mesons Into Higgs Particles*, *Phys. Rev. Lett.* **39** (1977) 1304 [[INSPIRE](#)].
- [60] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Remarks on Higgs Boson Interactions with Nucleons*, *Phys. Lett.* **78B** (1978) 443 [[INSPIRE](#)].
- [61] T. Inami, T. Kubota and Y. Okada, *Effective Gauge Theory and the Effect of Heavy Quarks in Higgs Boson Decays*, *Z. Phys. C* **18** (1983) 69 [[INSPIRE](#)].
- [62] NNPDF collaboration, *Parton distributions for the LHC Run II*, *JHEP* **04** (2015) 040 [[arXiv:1410.8849](#)] [[INSPIRE](#)].
- [63] A. Buckley et al., *LHAPDF6: parton density access in the LHC precision era*, *Eur. Phys. J. C* **75** (2015) 132 [[arXiv:1412.7420](#)] [[INSPIRE](#)].

- [64] X. Chen, J. Cruz-Martinez, T. Gehrmann, E.W.N. Glover and M. Jaquier, *NNLO QCD corrections to Higgs boson production at large transverse momentum*, *JHEP* **10** (2016) 066 [[arXiv:1607.08817](#)] [[INSPIRE](#)].
- [65] A. Gehrmann-De Ridder, T. Gehrmann, E.W.N. Glover, A. Huss and T.A. Morgan, *Precise QCD predictions for the production of a Z boson in association with a hadronic jet*, *Phys. Rev. Lett.* **117** (2016) 022001 [[arXiv:1507.02850](#)] [[INSPIRE](#)].
- [66] A. Gehrmann-De Ridder, T. Gehrmann, E.W.N. Glover, A. Huss and T.A. Morgan, *NNLO QCD corrections for Drell-Yan p_T^Z and ϕ^* observables at the LHC*, *JHEP* **11** (2016) 094 [Erratum *ibid.* **10** (2018) 126] [[arXiv:1610.01843](#)] [[INSPIRE](#)].
- [67] A. Gehrmann-De Ridder, T. Gehrmann, E.W.N. Glover, A. Huss and D.M. Walker, *Next-to-Next-to-Leading-Order QCD Corrections to the Transverse Momentum Distribution of Weak Gauge Bosons*, *Phys. Rev. Lett.* **120** (2018) 122001 [[arXiv:1712.07543](#)] [[INSPIRE](#)].
- [68] T. Gehrmann, M. Jaquier, E.W.N. Glover and A. Koukoutsakis, *Two-Loop QCD Corrections to the Helicity Amplitudes for $H \rightarrow 3$ partons*, *JHEP* **02** (2012) 056 [[arXiv:1112.3554](#)] [[INSPIRE](#)].
- [69] L.J. Dixon and Y. Sofianatos, *Analytic one-loop amplitudes for a Higgs boson plus four partons*, *JHEP* **08** (2009) 058 [[arXiv:0906.0008](#)] [[INSPIRE](#)].
- [70] S. Badger, E.W. Nigel Glover, P. Mastrolia and C. Williams, *One-loop Higgs plus four gluon amplitudes: Full analytic results*, *JHEP* **01** (2010) 036 [[arXiv:0909.4475](#)] [[INSPIRE](#)].
- [71] S. Badger, J.M. Campbell, R.K. Ellis and C. Williams, *Analytic results for the one-loop NMHV Hqgg amplitude*, *JHEP* **12** (2009) 035 [[arXiv:0910.4481](#)] [[INSPIRE](#)].
- [72] V. Del Duca, A. Frizzo and F. Maltoni, *Higgs boson production in association with three jets*, *JHEP* **05** (2004) 064 [[hep-ph/0404013](#)] [[INSPIRE](#)].
- [73] L.J. Dixon, E.W.N. Glover and V.V. Khoze, *MHV rules for Higgs plus multi-gluon amplitudes*, *JHEP* **12** (2004) 015 [[hep-th/0411092](#)] [[INSPIRE](#)].
- [74] S.D. Badger, E.W.N. Glover and V.V. Khoze, *MHV rules for Higgs plus multi-parton amplitudes*, *JHEP* **03** (2005) 023 [[hep-th/0412275](#)] [[INSPIRE](#)].
- [75] L. Cieri, in preparation.
- [76] L. Cieri, in preparation.
- [77] F. Dulat, A. Lazopoulos and B. Mistlberger, *iHixs 2 — Inclusive Higgs cross sections*, *Comput. Phys. Commun.* **233** (2018) 243 [[arXiv:1802.00827](#)] [[INSPIRE](#)].
- [78] C. Anastasiou et al., *High precision determination of the gluon fusion Higgs boson cross-section at the LHC*, *JHEP* **05** (2016) 058 [[arXiv:1602.00695](#)] [[INSPIRE](#)].
- [79] S. Catani, L. Cieri, G. Ferrera, D. de Florian and M. Grazzini, *Vector boson production at hadron colliders: a fully exclusive QCD calculation at NNLO*, *Phys. Rev. Lett.* **103** (2009) 082001 [[arXiv:0903.2120](#)] [[INSPIRE](#)].
- [80] S. Catani, L. Cieri, D. de Florian, G. Ferrera and M. Grazzini, *Diphoton production at hadron colliders: a fully-differential QCD calculation at NNLO*, *Phys. Rev. Lett.* **108** (2012) 072001 [Erratum *ibid.* **117** (2016) 089901] [[arXiv:1110.2375](#)] [[INSPIRE](#)].
- [81] M. Höschele, J. Hoff, A. Pak, M. Steinhauser and T. Ueda, *MT: A Mathematica package to compute convolutions*, *Comput. Phys. Commun.* **185** (2014) 528 [[arXiv:1307.6925](#)] [[INSPIRE](#)].

- [82] M. Höschele, J. Hoff, A. Pak, M. Steinhauser and T. Ueda, *Higgs boson production at the LHC: NNLO partonic cross sections through order ϵ and convolutions with splitting functions to N^3LO* , *Phys. Lett. B* **721** (2013) 244 [[arXiv:1211.6559](#)] [[INSPIRE](#)].
- [83] T. Gehrmann and E. Remiddi, *Two loop master integrals for $\gamma^* \rightarrow 3$ jets: The planar topologies*, *Nucl. Phys. B* **601** (2001) 248 [[hep-ph/0008287](#)] [[INSPIRE](#)].
- [84] D. Maître, *HPL, a mathematica implementation of the harmonic polylogarithms*, *Comput. Phys. Commun.* **174** (2006) 222 [[hep-ph/0507152](#)] [[INSPIRE](#)].
- [85] E. Remiddi and J.A.M. Vermaseren, *Harmonic polylogarithms*, *Int. J. Mod. Phys. A* **15** (2000) 725 [[hep-ph/9905237](#)] [[INSPIRE](#)].
- [86] A.B. Goncharov, *Multiple polylogarithms, cyclotomy and modular complexes*, *Math. Res. Lett.* **5** (1998) 497 [[arXiv:1105.2076](#)] [[INSPIRE](#)].
- [87] A.B. Goncharov, *Multiple polylogarithms and mixed Tate motives*, [math/0103059](#).
- [88] C. Duhr, H. Gangl and J.R. Rhodes, *From polygons and symbols to polylogarithmic functions*, *JHEP* **10** (2012) 075 [[arXiv:1110.0458](#)] [[INSPIRE](#)].
- [89] H. Frellesvig, D. Tommasini and C. Wever, *On the reduction of generalized polylogarithms to Li_n and $Li_{2,2}$ and on the evaluation thereof*, *JHEP* **03** (2016) 189 [[arXiv:1601.02649](#)] [[INSPIRE](#)].
- [90] C.W. Bauer, A. Frink and R. Kreckel, *Introduction to the GiNaC framework for symbolic computation within the C++ programming language*, *J. Symb. Comput.* **33** (2000) 1 [[cs/0004015](#)] [[INSPIRE](#)].
- [91] J. Vollinga and S. Weinzierl, *Numerical evaluation of multiple polylogarithms*, *Comput. Phys. Commun.* **167** (2005) 177 [[hep-ph/0410259](#)] [[INSPIRE](#)].